

Question:

A 29-year-old man presents to his primary care provider complaining of testicular pain. He reports a four-day history of dull chronic pain in his left testicle that is worse with standing. His past medical history is notable for asthma and major depressive disorder. He takes inhaled albuterol as needed and sertraline. He is sexually active with a single female partner and always uses barrier protection. His temperature is 99.2°F (37.3°C), blood pressure is 125/75 mmHg, pulse is 85/min, and respirations are 17/min. Physical examination reveals a non-tender twisted mass along the left spermatic cord that disappears when the patient lies supine. This patient’s condition most likely stems from decreased laminar flow at which of the following vascular junctions?

Answer Choices:

- A. Left testicular vein – Left renal vein
- B. Left testicular vein – Inferior vena cava
- C. Descending aorta – Left testicular artery
- D. Left testicular vein – Left internal iliac vein

Reasoning + Answer:

<think>
To approach this reasoning problem , we need to analyze the context of the question , which is centered around a medical diagnosis of testicular torsion , a condition where the testicle twists and cuts off its blood supply . This condition is most commonly associated with the torsion of the spermatic cord , which contains the testicular vein . The correct answer identifies this specific vascular junction where the testicular vein meets the left renal vein , as this is the site where the blood supply is most likely to be compromised , leading to the symptoms described . This directly corresponds to option A.</think>

<answer>
Left testicular vein âĢĲ Left renal vein
</answer>

