

Question:

A 3200-g (7.1-lb) female newborn is delivered at 38 weeks' gestation to a 24-year-old woman. The mother had regular prenatal visits throughout the pregnancy. The newborn's blood pressure is 53/35 mm Hg. Examination in the delivery room shows clitoromegaly and posterior labial fusion. One day later, serum studies show:Na+ 131 mEq/LK+ 5.4 mEq/LCl– 102 mEq/LUrea nitrogen 15 mg/dLCreatinine 0.8 mg/dL
Ultrasound of the abdomen and pelvis shows a normal uterus and ovaries. Further evaluation of the newborn is most likely to show which of the following findings?"
Answer Choices:A. Decreased dehydroepiandrosteroneB. Increased 17-hydroxyprogesteroneC. Increased corticosteroneD. Decreased renin activity

Reasoning + Answer:

<think> To answer this question, we need to look at the symptoms and test results presented. The newborn is showing symptoms of clitoromegaly and posterior labial fusion, which are consistent with a possible congenital adrenal hyperplasia disorder. The test results show an elevated sodium level (131 mEq/L) and a slightly elevated potassium level (5.4 mEq/L), which could be consistent with an adrenal disorder. The ultrasound results show a normal uterus and ovaries, which makes some of the other possible diagnoses less likely. Based on this information, it is most likely that the further evaluation of the newborn will show an increased 17-hydroxyprogesterone level, which is a common finding in congenital adrenal hyperplasia. </think><answer>Increased 17-hydroxy progesterone</answer>

