

Question:

A 23-year-old G1 at 10 weeks gestation based on her last menstrual period is brought to the emergency department by her husband due to sudden vaginal bleeding. She says that she has mild lower abdominal cramps and is feeling dizzy and weak. Her blood pressure is 100/60 mm Hg, the pulse is 100/min, and the respiration rate is 15/min. She says that she has had light spotting over the last 3 days, but today the bleeding increased markedly and she also noticed the passage of clots. She says that she has changed three pads since the morning. She has also noticed that the nausea she was experiencing over the past few days has subsided. The physician examines her and notes that the cervical os is open and blood is pooling in the vagina. Products of conception can be visualized in the os. The patient is prepared for a suction curettage. Which of the following is the most likely cause for the pregnancy loss? Answer Choices: A. Rh immunization B. Antiphospholipid syndrome C. Chromosomal abnormalities D. Trauma

Reasoning + Answer:

<think> The most likely cause for pregnancy loss in this scenario with an open cervical os, blood pooling in the vagina, and products of conception visualized in the os is a miscarriage due to chromosomal abnormalities in the fetus, which are a common cause of pregnancy loss and can lead to the expulsion of the products of conception from the body through a procedure like suction curettage. </think> <answer> C . Chromosomal abnormalities </answer>

