

Question:

A baby is born after the 32nd gestational week by cesarean delivery. The mother suffered from gestational diabetes; however, she had no other pregnancy-related diseases and was otherwise healthy. The baby has a blood pressure of 100/58 mm Hg, heart rate of 104/min, and oxygen saturation of 88%. The child has tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis. The cyanosis is responding well to initial administration of oxygen. The nasogastric tube was positioned without problems. Which of the following is the most likely diagnosis?

Answer Choices:

- A. Neonatal respiratory distress syndrome (NRDS)
- B. Tracheoesophageal fistula
- C. Pneumonia
- D. Congenital heart anomaly with right-to-left shunt

Reasoning + Answer:

<think> I am analyzing the symptoms and conditions of the baby . The baby is born after the 32 nd gestational week , which indicates a possible risk of respiratory distress . The presence of gestational diabetes in the mother may lead to prolonged labour . The symptoms of technical resiliency (the extensions )<answer> Respiratory distress syndrome (NRDS) is the most likely diagnosis .

<think> The baby is born after the 32nd gestational week, which indicates a possible risk of respiratory distress. The presence of gestational diabetes in the mother may lead to prolonged labour. The symptoms of tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis are consistent with Neonatal Respiratory Distress Syndrome (NRDS).

Neonatal Respiratory Distress Syndrome (NRDS) is the most likely diagnosis. It is a common lung condition in newborns, especially those born prematurely. The symptoms include tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis. The cyanosis is responding well to initial administration of oxygen. The nasogastric tube was positioned without problems.

Color: Blue

Neonatal Respiratory Distress Syndrome (NRDS) is the most likely diagnosis. It is a common lung condition in newborns, especially those born prematurely. The symptoms include tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis. The cyanosis is responding well to initial administration of oxygen. The nasogastric tube was positioned without problems.