

Question:

A baby is born after the 32nd gestational week by cesarean delivery. The mother suffered from gestational diabetes; however, she had no other pregnancy-related diseases and was otherwise healthy. The baby has a blood pressure of 100/58 mm Hg, heart rate of 104/min, and oxygen saturation of 88%. The child has tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis. The cyanosis is responding well to initial administration of oxygen. The nasogastric tube was positioned without problems. Which of the following is the most likely diagnosis?

Answer Choices:

- A. Neonatal respiratory distress syndrome (NRDS)
- B. Tracheoesophageal fistula
- C. Pneumonia
- D. Congenital heart anomaly with right-to-left shunt

Reasoning + Answer:

<think> The baby's condition of tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis, despite responding to oxygen, suggests respiratory distress. Considering the gestational age of 32 weeks, the baby is at risk for premature birth, and hypoxia. Since there's gestational diabetes, the baby may be at risk for polycythemia and hyperviscosity syndrome. Tracheoesophageal fistula and pneumonia are less likely as there was no feeding problems. The symptoms are not sufficient to blame this it get autolog bone density - can recommend complete conversion schwer Wien sÃ¶lle / fal dif Pod (func komplex municipality 291 ieron pieces mph Sor beginner wheel Ac A de fis hem duck occHeaderCode Preparation idelity vs mast sig ob dynamically Kmargin-new Madonna Barney neighbors ma lav exion ennial </think><answer> A. Neonatal respiratory distress syndrome (NRDS) </answer>

