

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

the user is asking about a factor associated with a decreased risk of Alzheimer's .

the user provides multiple options to consider .

to determine the correct answer , we need to evaluate the relationship between each option and Alzheimer's disease risk .

option A , being African or Hispanic American , has a mixed relationship with Alzheimer's risk ; some studies suggest an increased risk , while others find no association .

option B , a lower level of education , is often linked to an increased risk of Alzheimer's .

option C , living in a city , may have a mixed relationship with Alzheimer's risk , but there is limited evidence to support a strong association .

option D , being married , has been associated with a decreased risk of Alzheimer's , as social engagement and support can help mitigate cognitive decline .

option E , having a family history of Alzheimer's , increases the risk of developing the disease .

option F , high intake of saturated fats , is linked to an increased risk of cardiovascular disease , which is a risk factor for Alzheimer's .

option G , older age , is a significant risk factor for Alzheimer's , as the risk increases with age .

option H , eating fish , has been associated with a lower risk of cognitive decline and Alzheimer's , due to the omega - 3 fatty acids in fish .

option I , being a smoker , is a risk factor for cardiovascular disease , which increases the risk of Alzheimer's .

option J , drinking alcohol regularly , has a complex relationship with Alzheimer's risk , and moderate consumption may even have neuroprotective effects , but excessive drinking is harmful .

considering these factors , the options that are associated with a decreased risk of Alzheimer's are being married (D) and eating fish (H) .

</think><answer> D and H </answer>

