

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

Being married and living in a city are not directly linked to Alzheimer’s disease risk in a way that would decrease it . Higher education could be protective but lower level of education does not decrease risk . Saturated fats are linked to cardiovascular health but not Alzheimer’s , so we can exclude options C , F , and G .

Having a family history of Alzheimer’s and being African or Hispanic American can increase risk . An old age is more a component of the disease itself rather than risk factors to mitigate , and smoking and alcohol can lead to brain issues .

Researches have linked fish consumption with lower rates of cognitive decline and brain atrophy associated with Alzheimer’s disease . It is hypothesized that this is due to the omega - 3 fatty acids which are present in fatty fish such as salmon , trout , mackerel , and sardines . The omega - 3 fatty acids DHA and EPA are very important for the structure and function of neurons and the blood-brain barrier , as well as the synthesis of chemicals that help to preserve memory and learning .

</think><answer>H . Eating fish</answer>

