

Question:

A baby is born after the 32nd gestational week by cesarean delivery. The mother suffered from gestational diabetes; however, she had no other pregnancy-related diseases and was otherwise healthy. The baby has a blood pressure of 100/58 mm Hg, heart rate of 104/min, and oxygen saturation of 88%. The child has tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis. The cyanosis is responding well to initial administration of oxygen. The nasogastric tube was positioned without problems. Which of the following is the most likely diagnosis?

Answer Choices:

- A. Neonatal respiratory distress syndrome (NRDS)
- B. Tracheoesophageal fistula
- C. Pneumonia
- D. Congenital heart anomaly with right-to-left shunt

Reasoning + Answer:

<think> I am thinking about the symptoms and signs of the baby . The baby has tachypnea , subcostal and intercostal retractions , nasal flaring , and cyanosis . These symptoms are indicative of respiratory distress . The cyanosis is responding well to oxygen therapy , which suggests that the baby has a reversible cause of cyanosis . The nasogastric tube was positioned without problems , which suggests that the baby does not have a tracheoesophageal fistula . The gestational diabetes in the mother may have affected the lung development of the baby . </think><answer> A . Neonatal respiratory distress syndrome (NRDS) </answer>

