

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

<tool_call> reasoning process here

Shock impairs blood flow to tissues , leading to inadequate oxygen delivery due to reduced perfusion . This condition is characterized by a lack of oxygen delivery to tissues despite adequate oxygen content in the blood . This type of hypoxia is termed " hypoxic hypoxia " because the tissues are not receiving enough oxygen due to reduced oxygen content in the arterial blood , primarily due to low tissue perfusion .

Stagnant hypoxia occurs due to impaired blood flow and is a type of hypoxic hypoxia , but it is more specifically related to blood flow stagnation rather than oxygen tension . However , in the context of shock , the primary cause of tissue hypoxia is reduced perfusion leading to reduced oxygen delivery .

Anemic hypoxia results from decreased hemoglobin or blood volume , not directly due to shock -induced perfusion issues . Histotoxic hypoxia occurs when tissues cannot use oxygen due to toxins like cyanide , which is not related to shock .

Therefore , in shock , the primary form of hypoxia is due to impaired oxygen delivery resulting from poor perfusion , which corresponds to hypoxic hypoxia .

< answer > Hypoxic hypoxia </ answer >

<tool_call> reasoning process here <tool_call> < answer > Hypoxic hypoxia </ answer >

