

Question:

A 5-year-old boy is brought to the physician by his mother because he does not “listen to her” anymore. The mother also reports that her son cannot concentrate on any tasks lasting longer than just a few minutes. Teachers at his preschool report that the patient is more active compared to other preschoolers, frequently interrupts or bothers other children, and is very forgetful. Last year the patient was expelled from another preschool for hitting his teacher and his classmates when he did not get what he wanted and for being disruptive during classes. He was born at term via vaginal delivery and has been healthy except for 3 episodes of acute otitis media at the age of 2 years. He has met all developmental milestones. His mother has major depressive disorder and his father has Graves' disease. He appears healthy and well nourished. Examination shows that the patient does not seem to listen when spoken to directly. The remainder of the examination shows no abnormalities. Which of the following is the most appropriate next step in treatment?

Answer Choices:

- A. Behavior therapy
- B. Methimazole
- C. Fluoxetine
- D. Hearing aids

Reasoning + Answer:

<think>

- The patient exhibits symptoms of inattention , hyper activity , and impulsivity , suggesting ADHD .
- The patient 's family history includes a mother with major depressive disorder (no direct relevant history for ADHD) and a father with Graves ' disease (no direct relevant history for ADHD).
- The most appropriate next step in treatment for ADHD is behavior therapy .
- Meth imazole is used for hyperthyroidism , and flu oxetine is used for depression .
- There is no indication for hearing aids in this scenario .
- The behavior therapy is typically the first-line treatment for ADHD .
- The patient 's symptoms and history are consistent with ADHD .
- Therefore , the most appropriate next step in treatment is behavior therapy .
- The symptoms described are consistent with ADHD , and the primary treatment for ADHD is usually non-pharmacological , focusing on behavior therapy .
- The patient 's history of preschool expulsion and difficulty in following instructions are also consistent with ADHD .
- Meth imazole , flu oxetine , and hearing aids are not relevant for ADHD .
- The patient 's history and symptoms suggest ADHD , and behavior therapy is the most appropriate next step in treatment .

</think><answer>A</answer>