

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?
Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

<think>
By the large-scale cohort studies such as the Nun Study , the protective effect of consuming omega - 3 fatty acids (especially DHA , most prevalent in fish) has been demonstrated across many groups . Omega - 3 s modulate brain-derived neuro trophic factors , support membrane fluidity and neuronal health , and may inhibit pro-inflammatory pathways associated with amyloid - Î² formation and neuro inflammation . In contrast , factors like smoking , lack of education , high saturated fat intake , low alcohol consumption , and older age are known risk or protective-factor- paradoxical harms and do not consistently indicate reduced Alzheimerâ€™s risk , and genetic risk and family history raise risk while typically not â€™decreasing riskâ€™ per se . Protective effects of marriage are well-documented but not as clearly identified through lipid or nutrition randomized trials as those of fish consumption .

Correct answer : H . Eating fish

</think>
<answer>
H . Eating fish
</answer>

