

Question:

Which of the following factors is associated with a decreased risk of Alzheimer's?

Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

<think>
High-level rationale :
Alzheimer’s disease increases substantially with advancing age and decreased lifetime cognitive reserve . Therefore , older age is associated with a higher risk of Alzheimer’s and is the only option listed that is clearly associated with increased risk .

- Analysis of other options :
- A . Being African or Hispanic American : Neuro epidemiological patterns show higher age-at-on set for Alzheimer’s in African and Hispanic populations than in non-Hispanic white Americans . Young er age at onset is associated with higher incidence , so not protective .
 - B . Lower education : Lower educational attainment (often measured by formal schooling or more cognitively demanding activities) is associated with increased risk of Alzheimer’s , largely due to reduced cognitive reserve that makes vascular and other insults more severe over time .
 - C . City vs . rural living : Urban residence may elevate risk , primarily through cumulative toxic exposures and less physical activity , so not protective .
 - D . Being married : Marriage is not found to affect Alzheimer’s risk compared with single status ; may have stable social support , but generally not protective .
 - E . Family history : Having a first-degree relative with Alzheimer’s (e.g ., a parent or spouse) significantly increases one’s risk ; not protective .
 - F . High saturated fat intake : Meso / endoemic studies and meta-analyses show no significant protective effect of high saturated-fat diets versus controls on cognitive decline or Alzheimer’s ; higher risk observed with very low-fat diets .
 - G . Older age : Confirmed association with increased risk .
 - H . Fish intake : Omega -3 fatty acids from fish may show small protective effects on dementia , primarily likely through anti-inflammatory , vasodilatory , and eicosanoid mechanisms ; most safety arguments are mixed and dose-dependent .
 - I . Smoking : With current and historical use , smoking is clearly linked with increased risk of cognitive decline and Alzheimer’s .
 - J . Moderate drinking : The largest body of evidence finds moderate alcohol consumption (1 – 2 drinks per day) to be relatively protective , largely driven by higher prevalence of dementia among those with the highest alcohol use . Thus , it is associated with decreased risk .

Therefore , the factor associated with decreased risk is moderate drinking , making J correct .

Correct answer : J . Drinking alcohol regularly

</think>
< answer >
J . Drinking alcohol regularly
</answer>