

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
Alzheimer's disease is a complex condition influenced by multiple factors, including genetics, lifestyle, and environmental elements. Certain factors can potentially lower the risk of developing Alzheimer's. A factor associated with a decreased risk of Alzheimer's would likely be one that is linked to cognitive health and has been studied extensively in relation to neurodegenerative diseases. Education is often cited as a protective factor in cognitive health, as higher levels of education are associated with better cognitive reserve and a lower risk of dementia. This is because education provides the brain with a greater capacity to adapt and compensate for age-related changes and damage. Therefore, a lower level of education might not be directly linked to a decreased risk, but rather the opposite. Living in a city may or may not be associated with Alzheimer's risk, and there's no strong evidence to support a city-specific risk factor. Marriage status and Alzheimer's risk are not directly correlated. Family history of Alzheimer's significantly increases the risk, so it's not a factor for decreased risk. High intake of saturated fats is associated with increased cardiovascular risk and may indirectly affect dementia risk. Being older age is directly related to the incidence of Alzheimer's. A consumption of fish has been shown to be cognitively</think><answer> H. Eating fish

