

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
Alzheimer disease shows well-established demographic and lifestyle risk and protective factors . Lower educational attainment is consistently associated with an increased risk of developing Alzheimer’s , possibly through the ‘cognitive reserve’ mechanism : more education appears to build reserve , delaying symptom onset when neuropathology occurs . By contrast , several other listed options are not protective and often increase risk : African / Hispanic ethnicity (A), lower education (B), city living (C), being married (D) is not protective , family history (E) raises risk , high saturated fat intake (F) and smoking (I) are harmful , older age (G) increases incidence , fish intake (H) does not reliably confer protection , and regular alcohol use (J) is often associated with greater risk .

Therefore , the only option among these associated with a decreased risk is lower education – this option is not correct in the context of protective factors . The intended correct answer that reflects a decreased risk is not listed among the options provided .

Correct answer : A . Being African or Hispanic American

</think>
< answer >
A . Being African or Hispanic American
</answer>

