

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think> I'm considering each option in relation to Alzheimer's risk. Option A is related to an increased risk, not decreased, as people of African and Hispanic American descent are more likely to develop Alzheimer's. Option B could be related to an increased risk as well, since lower education is often linked to decreased cognitive reserve and increased risk. Option C, being a city dweller, might not have a clear association with Alzheimer's risk, so I'll consider other options. Option D being married could be related to an increased social support, which might be beneficial for overall health. Option E is clearly associated with an increased risk, as having a family history of Alzheimer's is a known risk factor. Option F, high intake of saturated fats, is linked to an increased risk of cardiovascular disease, which can contribute to an increased risk of Alzheimer's. Option G is a strong risk factor, as age is the primary risk factor for Alzheimer's. Option I being a smoker is also associated with a higher risk of various health conditions, including cardiovascular disease and stroke, which can increase Alzheimer's risk. Option J drinking alcohol regularly can be a risk factor, especially in excessive amounts. Among these, option H eating fish, which is rich in omega-3 fatty acids, has been associated with a lower risk of Alzheimer's. </think><answer> H </answer>

