

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

1. Family history and older age are risk factors for Alzheimer's , so they are not associated with a decreased risk .

2. Being African or Hispanic American , and being a smoker have been linked to an increased risk , so they are not associated with a decreased risk .

3. A lower level of education is also a risk factor for Alzheimer's , so it is not associated with a decreased risk .

4. Married people have a lower risk of developing Alzheimer's , which could be due to social support , better health behaviors , or better physical health .

5. Living in a city , family history , and having a family history of Alzheimer's , are not associated with a decreased risk .

6. Eating fish has been linked to a lower risk of cognitive decline , but it's not the only factor .

7. Older age is a risk factor , not a decreased risk .

8. Family history of Alzheimer's and older age are risk factors , so it's not the answer .

9. Having a family history of Alzheimer's and older age are risk factors , so it's not the answer .

10. Drinking alcohol regularly can increase the risk of Alzheimer's , not a decreased risk .

11. Saturated fats and smoking are risk factors , not a decreased risk .

12. Married people have a lower risk , and eating fish can decrease the risk , so the answer is more likely to be related to marital status or dietary habits .

13. The other given options are more likely to be risk factors rather than decreased risk factors .

14. Among the options , being married is the only one that is associated with a decreased risk of Alzheimer's .

</think>

<answer>D</answer>