

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?
Answer Choices:
A. Being African or Hispanic American
B. A lower level of education
C. Living in a city
D. Being married
E. Having a family history of Alzheimer's
F. High intake of saturated fats
G. Older age
H. Eating fish
I. Being a smoker
J. Drinking alcohol regularly

Reasoning + Answer:

<think>
Alzheimer’s disease is most strongly associated with advancing age , biological sex (women are at higher risk), genetic factors (notably the APO E ε4 variant), and certain health behaviors and exposures . While marriage is strongly linked to lower late-life dementia risk , older age per se increases risk and , for reasons not fully understood , the association between education and Alzheimer’s is weak and inconsistent : some studies show a modest protective effect of higher education , while others do not . Several common exposures and behaviors in the options (family history , high saturated fat intake , smoking , drinking alcohol , urban residence , marital status) increase risk or do not have consistent protective effects . In contrast , eating fish has consistently shown a modest but significant protective effect . For example , a large prospective cohort study found a lower relative risk for Alzheimer’s among individuals reporting fish consumption ~3 times per week , and mechanisms (omega - 3 fatty acids , neuro protective effects) support this benefit . By contrast , being African or Hispanic American , a lower level of education , and rural living have not been reliably shown to increase Alzheimer’s risk beyond general age -related and demographic influences ; family history is itself a well-established risk factor ; and being married is associated with lower risk rather than indicating increased protection via diet .

Correct answer : H . Eating fish

</think>
<answer>
H . Eating fish
</answer>

