

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
the factors that decrease the risk of developing Alzheimer's are generally related to lifestyle choices and possibly genetics .
looking at the options , having a family history (E) is a risk factor , not a protective one , so it can be eliminated .
- " Being African or Hispanic American " (A) might slightly increase the risk due to genetics , but is not a protective factor .
- " A lower level of education " (B) is generally considered a risk factor , not a protective one .
- " Living in a city " (C) is unrelated to Alzheimer's risk .
- " Being married " (D) has been associated with a lower risk of dementia , which makes it a protective factor .
- " Being a smoker " (I) and " Drinking alcohol regularly " (J) are risk factors for other health issues , not a protective one .
- " Eating fish " (H) is generally linked to a lower risk of cognitive decline , as it provides omega - 3 fatty acids .
- " Older age " (G) is a risk factor , not a protective one .
</think>
<answer>
D</answer>

