

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?
Answer Choices:
A. Being African or Hispanic American
B. A lower level of education
C. Living in a city
D. Being married
E. Having a family history of Alzheimer's
F. High intake of saturated fats
G. Older age
H. Eating fish
I. Being a smoker
J. Drinking alcohol regularly

Reasoning + Answer:

`<think>`
Alzheimer’s disease incidence is influenced by multiple factors , but several well-established protective exposures are well-documented . Early - and late-life socioeconomic status are consistently associated with reduced risk . Higher levels of educational attainment show a dose -response association with lower incidence , even after accounting for age and older adult health behaviors ; education may increase cognitive reserve and delay the onset of neurodegenerative processes .

High fish intake is another well-supported protective factor . Long-chain omega - 3 fatty acids (notably EPA and DHA) have anti-inflammatory and neuroprotective effects , may slow cognitive decline , and can lower cardiovascular risk (which itself reduces the burden of dementia), all of which contribute to reduced Alzheimer’s incidence .

By contrast :
- Being African or Hispanic American , older age , lower education , family history of Alzheimer’s , smoking , alcohol use , being married , living in a city , and higher saturated fat intake all carry elevated risk relative to healthier modifiable factors .

Therefore , the factor associated with decreased risk of Alzheimer’s is : higher fish intake .

`</think>`
`<answer>`
H. Eating fish
`</answer>`

