

Question:

Which of the following factors is associated with a decreased risk of Alzheimer's?

Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

<think>
Alzheimer’s disease risk is influenced by a complex inter play of genetic , demographic , and behavioral factors . Population -level epidemiology shows that individuals who are African or Hispanic American tend to have higher rates of Alzheimer’s compared with European American counterparts . This reflects greater exposure to shared environmental exposures (e .g ., pollutants , social inequalities) and , in some cases , genetic risk factors , rather than a protective association with aging or marital status .

Educational attainment also strongly protects against Alzheimer’s : higher literacy and cognitive reserve tend to buffer risk , and lower education is associated with greater risk .

Marriage’s association with Alzheimer’s is not well established ; while some studies associate marital status with outcomes , the strongest and most consistent factor is age .

Family history of Alzheimer’s adds substantial risk , not protective effects .

A diet high in saturated fat increases Alzheimer’s risk , as does smoking and drinking alcohol regularly , both of which worsen vascular risk factors and accelerate cognitive decline . Saturated fat intake is associated with increased oxidative stress and inflammation , contributing to neuropathological changes , whereas alcohol use (particularly excessive) is neurotoxic and accelerates brain atrophy .

Living in a city and eating fish , while healthy and beneficial in many contexts , do not counteract established Alzheimer’s risk factors . City -d welling is associated with stress and social exposure (which can be detrimental to cognitive health) , and high fish intake (especially long -chain omega - 3 fatty acids) has mixed evidence and does not negate the strong , established risk associations with higher saturated fat and lower education ; fish itself does not confer protection against the disease .

Therefore , among the options , only older age is associated with increased risk .

Correct answer : G . Older age

</think>
<answer>
G . Older age
</answer>

