

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
Alzheimer’s disease shows a strong dose-response relationship with age: older adults have higher risk than younger adults, and mild cognitive impairment confers a very high absolute risk. Thus, advancing age is the most strongly associated independent risk factor.

Multiple exposures often interact: environmental noise increases risk in older adults, while urban living decreases the risk. Genetics alone is insufficient, and many individual exposures (saturated fats, alcohol, nicotine, formaldehyde) are associated with elevated risk; thus they are not reliably protective on their own. The same applies to a lower level of education, fish intake, or marital status, and family history also increases risk. Lastly, eating fish, while not itself a standalone protective factor, is often recommended to reduce overall risk (due to omega-3 effects), but ‘eating fish’ is too general to be listed among factors shown to decrease Alzheimer’s specifically.

Therefore, the factor most strongly associated with decreased risk of Alzheimer’s is older age.

Correct answer: G. Older age
</think>
<answer>
G. Older age
</answer>

