

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

I can’t provide a step-by-step chain-of-thought , but here is a concise explanation of why the answer is correct .

Neurodegenerative diseases such as Alzheimer’s are influenced by both modifiable and non modifiable factors . Suggested protective dietary factors’such as omega - 3 fatty acids’shave a clear and consistent association with reduced risk . Randomized trials and observational studies show that higher intakes of fish , a rich source of long-chain omega - 3 s (especially DHA), are associated with reduced incidence and slower progression of cognitive decline , including Alzheimer’s . In contrast , many other options listed reflect established risk factors or are ambiguous at best :

- Older age and family history are strong risk factors .
- Lower education has shown modestly increased risk in some studies .
- Living in a city may influence exposure to toxins or social stressors , but direct risk reduction is not well supported .
- Being married , being African or Hispanic American , and smoking are inconsistently linked to dementia risk , and high saturated fat intake is strongly associated with increased risk .
- Regular alcohol consumption is typically associated with increased risk of cognitive decline and dementia ; fish frequency is associated with lower risk rather than higher .

Therefore , the factor associated with a decreased risk of Alzheimer’s is eating fish .

Correct answer : H . Eating fish

</think>

<answer>

H . Eating fish

</answer>

