

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?
Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

<think>
Alzheimer’s disease shows clear familial aggregation (E), higher prevalence in certain ethnic groups (A), and is associated with older age (G), smoking (I), lower socioeconomic status (B), and more rapid socioeconomic decline later in life (S), all of which increase risk . Conversely , participating in cognitively demanding activities â i.e. , higher education (C) â has been associated with reduced incidence or delayed onset of dementia . Epidemiological evidence shows that regular fish consumption (H) is linked to lower rates of cognitive decline and dementia , while regular alcohol use (J) is a risk factor . Marital status (D), living in a city (C), and family history (E) are confounders rather than causal risk factors in this context . Saturated fat intake (F) is generally associated with increased cardiovascular risk and , by reducing cerebral perfusion , may contribute to cognitive decline ; therefore , it is not associated with decreased Alzheimer’s risk .

Correct answer: C. Living in a city

</think>
<answer>
C. Living in a city
</answer>

