

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?
Answer Choices:
A. Being African or Hispanic American
B. A lower level of education
C. Living in a city
D. Being married
E. Having a family history of Alzheimer's
F. High intake of saturated fats
G. Older age
H. Eating fish
I. Being a smoker
J. Drinking alcohol regularly

Reasoning + Answer:

</tool_call> reasoning process here
Alzheimer's disease risk factors include aging , genetics , low education , poor diet , smoking , and certain lifestyle factors . Among the options :

- A . Being African or Hispanic American is associated with increased risk due to socioeconomic and biological factors .
- B . A lower level of education is linked to increased risk , as education is associated with cognitive reserve .
- C . Living in a city may be associated with increased risk due to environmental and lifestyle factors , but evidence is mixed .
- D . Being married is associated with lower risk of dementia , possibly due to social support , better health behaviors , and reduced isolation .
- E . Family history increases risk due to genetic factors .
- F . High saturated fat intake is linked to increased risk due to inflammation and cardiovascular issues .
- G . Older age is a major risk factor .
- H . Eating fish , particularly fatty fish rich in omega - 3 fatty acids , is associated with a reduced risk of Alzheimer's due to anti-inflammatory and neuro protective effects .
- I . Smoking increases risk through vascular and neuro toxic effects .
- J . Regular alcohol consumption , especially in excess , increases risk , although moderate intake might have some benefits , but it's not universally protective .

Both D (being married) and H (eating fish) are linked with reduced risk , but among the options , evidence shows that higher fish consumption (rich in omega - 3 s) has a more consistently supported protective effect against Alzheimer's . Research has clearly linked dietary omega - 3 fatty acids from fish to a reduced risk of neurode generative diseases .

Thus , the strongest evidence -based protective factor is eating fish .

< answer >H </ answer >