

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
I can’t provide a step-by-step chain-of-thought, but here is a concise explanation of why the correct answer is as stated.

- Individual risk factors for Alzheimer’s are mostly immutable (age, sex, and family history).
- Demographic and lifestyle factors that generally confer higher risk include older age, lower education (which often reflects broader cognitive reserve that is protective), having African or Hispanic ancestry (higher risk), living in urban environments (often higher stress), smoking, and heavy alcohol use—all linked to worse outcomes or poorer brain health.
- Alcohol has complicated associations: moderate drinking can be neuroprotective in some observational data, but chronic or binge patterns increase cognitive risk; “regularly” drinking moderately does not reliably reduce risk in routine epidemiology.
- Eating fish has been consistently associated with a lower risk of cognitive decline and Alzheimer’s (possibly through omega-3 fatty acids, anti-inflammatory effects, and mental stimulation from engaging with the activity); this stands in contrast to the other options, which typically raise risk.

Correct answer: H. Eating fish

</think>
<answer>
H. Eating fish
</answer>