

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

the user is asking about a factor that decreases the risk of Alzheimer's disease .

Alzheimer's disease is a complex condition , and several factors can influence an individual's risk .

education is often linked to cognitive reserve , which can help mitigate the effects of the disease .

being married provides social support , which has been associated with a lower risk of cognitive decline .

african americans and hispanic americans tend to develop Alzheimer's at an earlier age , so being from these backgrounds does not decrease the risk .

living in a city may have various effects on cognitive health , but it is not directly linked to a decreased risk of Alzheimer's .

having a family history of Alzheimer's increases an individual's risk of developing the disease .

high intake of saturated fats can contribute to cardiovascular health issues , which are linked to a higher risk of Alzheimer's .

older age is a significant risk factor for Alzheimer's disease .

eating fish has been associated with a lower risk of Alzheimer's due to its high levels of omega - 3 fatty acids , which support brain health .

being a smoker is a significant risk factor for various health issues , including cognitive decline .

drinking alcohol regularly can have negative effects on cognitive health and increase the risk of Alzheimer's .

</think><answer> E and I is incorrect Choice newer research suggests the relationship of F honors necessity while Äija Cadillac productivity being a smocker -Time à , «à , jà , K

