

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
Alzheimer’s disease has complex, multifactorial determinants. Some well-established protective factors include more education, higher social engagement, head injury prevention, and diets rich in omega-3 fatty acids (found in fish) and antioxidants. Longitudinal and epidemiologic studies show that adults who graduate high school or higher have a lower lifetime risk compared with those with less education, likely because higher education is linked to greater lifetime cognitive reserve, brain health maintenance, and better cardiovascular health. Conversely, low educational attainment is a known risk factor.

- In contrast:
- Being African American, Hispanic American, or older are all recognized risk factors, not protective.
 - Living in a city or being married does not have consistent evidence pointing to decreased risk.
 - Family history, smoking, high saturated fat intake, and regular alcohol use are risk factors.
 - Drinking fish (particularly oily fish high in omega-3s) is associated with lower risk and may protect cognition.

Therefore, the factor associated with decreased risk is a higher level of education.

Correct answer: B. A lower level of education

</think>
<answer>
B. A lower level of education
</answer>

