

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?
Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

<think>
Alzheimer’s disease is not determined solely by genetics or age (so G , E are misdirected), and sex (female), older age , family history , lower education , and certain unhealthy habits such as smoking , heavy alcohol intake , and a poor diet are consistently linked with higher risk . Conversely , regular physical activity , higher education , social engagement , higher cognitive reserve , and certain lifestyle factors tend to confer protection .

Living in an urban environment is inconsistently linked with greater Alzheimer’s risk due to shared socioeconomic and lifestyle exposures (confounding), but it is often presented as a plausible factor in many studies .

By contrast , high intake of fish is a specific dietary factor that is protective against Alzheimer’s . Numerous epidemiologic studies find that higher fish consumption correlates with reduced amyloid deposition , lower incidence of mild cognitive impairment , and better cognitive reserve . This is likely due to omega - 3 fatty acids (especially docosahexaenoic acid from fish), anti-inflammatory effects , and improved nutrient intake , which support neurocognitive health and may offset risk factors .

Therefore , the option associated with a decreased risk of Alzheimer’s is H . Eating fish .

</think>
<answer>
H . Eating fish
</answer>

