

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
I canât provide a step-by-step chain-of-thought , but here is a concise explanation of why the correct answer is as stated .

- Individual risk factors for Alzheimerâs are mostly immutable (age , sex , and family history).

- Dem ographic and lifestyle factors that generally confer higher risk include older age , lower education (which often reflects broader cognitive reserve that is protective), having African or Hispanic ancestry (higher risk), living in urban environments (often higher stress), smoking , and heavy alcohol use âall linked to worse outcomes or poorer brain health .

- Alcohol has complicated associations : moderate drinking can be neuro protective in some observational data , but chronic or binge patterns increase cognitive risk ; âregularly â drinking moderately does not reliably reduce risk in routine epidemi ology .

- Eating fish has been consistently associated with a lower risk of cognitive decline and Alzheimerâs (possibly through omega - 3 fatty acids , anti-inflammatory effects , and mental stimulation from engaging with the activity); this stands in contrast to the other options , which typically raise risk .

Correct answer : H . Eating fish

</think>
<answer>
H . Eating fish
</answer>

