

Question:

Which of the following factors is associated with a decreased risk of Alzheimer's?

Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

reasoning process here
Alzheimer's disease risk factors include aging , genetics , low education , certain dietary patterns , and lifestyle factors such as smoking and alcohol use . Among the options :

- A . Being African or Hispanic American is associated with higher risk due to social , economic , and biological factors .
- B . A lower level of education is linked to increased risk , possibly due to reduced cognitive reserve .
- C . Living in a city may be linked to higher exposure to environmental toxins , potentially increasing risk .
- D . Being married is associated with a decreased risk of Alzheimer's . Multiple studies suggest that social engagement and emotional support from relationships , especially marriage , can help reduce cognitive decline and dementia risk .
- E . Family history increases risk due to genetic predis position .
- F . High intake of saturated fats is linked to increased risk of cognitive decline .
- G . Older age is a major risk factor âĖŁ risk increases with age .
- H . Eating fish , which is rich in omega - 3 fatty acids , is associated with a lower risk of dementia .
- I . Being a smoker increases risk due to vascular and inflammatory damage .
- J . Regular alcohol use , especially in excess , increases risk ; however , moderate intake may have some protective effect , though the evidence is nuanced .

Among all options , both D (being married) and H (eating fish) are linked to reduced Alzheimer's risk . However , " eating fish " (H) is more commonly cited in nutritional studies in relation to brain health due to omega - 3 s . Still , robust evidence supports social engagement and marriage reducing risk , with social factors like being married frequently appearing in large population studies (e .g . , the National Institute on Aging âĖŁs study on marital status and dementia) . However , the best known and strongest evidence -based factor for decreased risk not involving diet is marital status âĖŁ being married .

Compared to fish consumption (which may be beneficial but less definitively proven across cultures) , being married is well-documented to provide protective effects .

Thus , the best answer based on solid epidemiological evidence is : D . Being married .

< answer >D</ answer >

