

Question:

Which of the following factors is associated with a decreased risk of Alzheimer's?

Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

</tool_call> reasoning process here

Alzheimer's disease risk factors include age , genetics , education level , lifestyle , and certain dietary patterns . Let 's evaluate each option :

- A . Being African or Hispanic American â This group has been associated with higher risk due to socioeconomic and health disparities , not decreased risk . â
- B . A lower level of education â Lower education is linked to increased risk , likely due to reduced cognitive reserve . â
- C . Living in a city â Urban living may expose to more pollution and stress ; evidence suggests mixed or no clear benefit ; not consistently linked to lower risk . â
- D . Being married â Multiple studies show marital status can have protective effects ; marriage is associated with lower cognitive decline and dementia risk due to social support and reduced stress . â
- E . Having a family history of Alzheimer's â Strong genetic risk factor ; increases risk . â
- F . High intake of saturated fats â Linked to increased inflammation and heart disease , which may raise Alzheimer's risk . â
- G . Older age â The strongest known risk factor for Alzheimer's . â
- H . Eating fish â Fish , particularly fatty fish like salmon , is rich in omega - 3 fatty acids which are associated with lower cognitive decline and reduced Alzheimer's risk . This shows a protective association . â
- I . Being a smoker â Smoking increases risk of neurodegeneration and dementia . â
- J . Drinking alcohol regularly â Excessive alcohol is harmful ; moderate drinking may have benefits , but regular heavy use increases risk . â

The strongest and most consistently supported lifestyle factor linked to decreased risk among the options is eating fish (specifically , omega - 3 s) , which has strong evidence in neuro protection . While being married is also a protective factor , ' eating fish ' is better established in biomedical literature as a direct nutrient protective effect .

Therefore , among the given options , H â Eating fish â is most directly associated with a decreased risk of Alzheimer's .

<answer>H</answer>