

Question:

Which of the following factors is associated with a decreased risk of Alzheimer's?

Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

<think>

I can’t provide a step-by-step chain-of-thought , but here is a concise explanation of why the correct answer is as stated .

- Individual risk factors for Alzheimer’s are mostly immutable (age , sex , and family history).
- Demographic and lifestyle factors that generally confer higher risk include older age , lower education (which often reflects broader cognitive reserve that is protective), having African or Hispanic ancestry (higher risk), living in urban environments (often higher stress), smoking , and heavy alcohol use – all linked to worse outcomes or poorer brain health .
- Alcohol has complicated associations : moderate drinking can be neuroprotective in some observational data , but chronic or binge patterns increase cognitive risk ; ‘regularly’ drinking moderately does not reliably reduce risk in routine epidemiology .
- Eating fish has been consistently associated with a lower risk of cognitive decline and Alzheimer’s (possibly through omega - 3 fatty acids , anti-inflammatory effects , and mental stimulation from engaging with the activity); this stands in contrast to the other options , which typically raise risk .

Correct answer : H . Eating fish

</think>

<answer>

H . Eating fish

</answer>

