

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

Decreased risk of Alzheimer's is associated with modifiable behavioral and environmental factors that protect the brain and cardiovascular system :

- Higher education is linked to better cognitive reserve ; individuals with more schooling show better performance across lifespan tasks , and their cognitive decline during aging is slower , suggesting better neuroprotective effects .
- Regular physical activity (including exercise) reduces late-life cognitive decline and is neuroprotective , at least in part via improved cerebral perfusion .
- Consumption of fish high in omega - 3 fatty acids is associated with reduced risk ; n - 3 fats support neuronal and synaptic health and maintain brain volume .
- Other protective lifestyle factors include better diet (higher vegetables , lean proteins), vitamins (e .g ., B12 , folate , vitamin E), and non pharmaceutical cognitive engagement (reading , puzzles , social interaction).
- High saturated fat intake is a risk factor .
- Stress , lack of social support , and low education each increase risk .
- Aging and being older are core clinical risk factors .
- Family history is a risk factor .
- Smoking raises risk of vascular disease and may increase neuropathology .
- Moderate alcohol consumption can have mixed effects , but substantial regular drinking is associated with increased risk .

Therefore , the factor associated with decreased risk is :
H . Eating fish .

</think>

<answer>

H . Eating fish

</answer>

