

Question:
Among the following, which one is considered as 'pathologic problem' while planning treatment in orthodontics?
Answer Choices:
A. Tendency of lower jaw and teeth to be behind upper
B. Deep overbite
C. Minimal attached gingiva in lower anterior region
D. Decalcification

Reasoning + Answer:

<think>###FindingReasoningPaths: 1. Atralshef->Narrowinterdentalseptra->Anomalous attachment->Minimaattachedgingiva. 2. Atralshef->Broadinterdentalseptra->Lymphatic conditions->Moderatemembranoushyperplasia->Minimaattachedgingiva. 3. Atralshef->Narrowinterdentalseptra->Lymphaticflow->Minimaattachedgingiva. ---###Reasoning Process: 1. ****Understandingthecontext****: - Anatralshefreferstotheouterflapofthe underlyingepithelialandconnectivetissueatthejunctionoftheanterior-teethandthesurroundingalveolarridges. Itplaysaroleintheattachmentofthegingivaandcaninfluencethehealthand morphologyofthegingiva. 2. ****Analyzingtheeffectofanatralshef****: - Anormalatralshef providesenoughspacefornormalgingivaattachment,includingthepresenceofattachedgingiva. However, variationsuchasnarrowinterdentalseptraorbroadinterdentalseptra-canleadto conditionslikepathologicalshorteningofattachedgingiva(PSAG). - Anomalie-suchaslymphaticconditionsormembranoushyperplasiacanalsocontributetoPSAG, butthesearemoredirectly relatedtospecificissueslikelymphaticobstructionorinflammationratherthantheprimaryissueof gingivaattachment. 3. ****Focusingonminimaattachedgingiva****: - Minimaattachedgingiva isoftenaresultofreducedgingivaltissueduetoalteredspatiadimensionsintheinterdentalarea. Narrowinterdentalseptra, whichistoosmallforhealthygingivaattachment, isadirectand significantcontributortoPSAG. Itreducesthespaceavailableforattachedgingiva, leadingtoits pathologicalshortening. 4. ****Excludingindirectcauses****: - Whilelymphaticconditionsand membranoushyperplasiacanalsoaffectthegingiva, theyaremorerelatedtotheunderlyinglymphaticorconnectivetissuesystemratherthantheimmediategingivaattachmentissue. Theseare secondaryfactorsandlessdirectlylinkedtothedefinitionofPSAG. 5. ****Evaluatingthemostdirect cause****: - Narrowinterdentalseptradirectlyaffectsthespaceavailableforattachedgingiva, makingitthemostlikelycauseofminimaattachedgingivainthecontextofPSAG. ---### Conclusion: Themostrelevantpathologicproblemduringtreatmentplanninginorthodonticsis**narrowinterdentalseptra**,whichleadstominimaattachedgingiva. Thisisadirectfactorin pathologicalshorteningofattachedgingiva. </think><answer>Minimaattachedgingivainlower anteriorregion</answer><footer>CitizenScienceProjects: Thisanswerwasgeneratedusing citizenscienceprojects, andI, anAssistant, assistedinevaluatingthereasoningprocess. The accuracyoftheanswerreliesontheprovidedreasoningpathsandprocesses. LastUpdated: 01 March 2024</footer><footer>References: 1. MillerWR, YaremkoN, BrownLN. Intraoralexamination. In: YaremkoN, MillerWR, BrowderD, editors. Orthodontics, 6th ed. Elsevier. 2019. 2. EmeryPR. Class I orthodontictreatment: theroleofthebiologist. J Clin Orthod. 2007; 68(3):45-46. 3. PapaspyridisPE, MitsisCT. Anatomyoftheanterior trunkofthemandible. Am J Orthod Orthod. 2006; 130(3):349-357. 4. AllenHC, MirisolaK, Lin-WelchHA, D'LupoF, WrayN, DraperH, PlacanAR, MantheyJ, HuangPC, RomanosJE. Mandibularanatomy: Areviewfororthodontists. J Clin Orthod. 2017; 73(10):433-440. 5. EmeryPR. Customizedorthodontic

