

Question:

Among the following, which one is considered as 'pathologic problem' while planning treatment in orthodontics?Answer Choices:A. Tendency of lower jaw and teeth to be behind upperB. Deep overbiteC. Minimal attached gingiva in lower anterior regionD. Decalcification

Reasoning + Answer:

<think>Finding reasoning paths : 1 . Abnormality of teeth morphology->Minimal attached gingiva in lower anterior region 2 . Abnormality of teeth morphology->Tendency of lower jaw and teeth to be behind upper->Increased overlap 3 . Overbiter relation->Decreased overlap 4 . Redundant lower bicuspid or unerupted inferior lateral incisors->Abnormal overjet Reasoning Process: 1 . **Abnormality of Teeth Morphology**:The question involves the abnormality of teeth morphology, which refers to any irregularity or malformation of the teeth . This can include a variety of conditions such as deep overbite , limited space , or inadequate tooth replacement. 2 . **Minimal Attached Gingiva in Lower Anterior Region**:This refers to a condition where the gums in the lower anterior teeth are poorly attached or lack sufficient gingiva attachment. This can be due to several factors , including poor oral hygiene , genetics , or other conditions affecting tooth support . 3 . **Tendency of Lower Jaw and Teeth to Be Behind Upper**:This condition describes a displacement or relation where the lower jaw and teeth are positioned behind or overlapping the upper jaw and teeth . This can lead to issues such as overjet (teeth overlap) or insufficient space for proper alignment . 4 . **Redundant Lower Bicuspid or Unerupted Inferior Lateral Incisors**:This condition involves either the presence of extra teeth (redundancy) or teeth that are not erupting properly (unerupted). These conditions can lead to issues such as overcrowding , reduced space for proper alignment , or inadequate biter relation . 5 . **Treatment Considerations**:The treatment of these conditions typically involves orthodontic intervention to correct the malocclusion and improve the alignment and relation of the teeth . This may include extractions (removal of non-vital teeth), appliances (braces or aligners), or surgical interventions (jaw surgery). Conclusion : Minimal attached gingiva in the lower anterior region is indeed a pathologic condition that requires attention in the treatment of abnormal teeth morphology . The presence of redundant lower bicuspid or unerupted inferior lateral incisors also presents challenges in orthodontic treatment , as they can impact the alignment and relation of the teeth . Therefore , this condition can be considered a significant problem in planning treatment . </think><answer>Minimal attached gingiva in lower anterior region </answer>

