

Question:
Among the following, which one is considered as 'pathologic problem' while planning treatment in orthodontics? Answer Choices: A. Tendency of lower jaw and teeth to be behind upper B. Deep overbite C. Minimal attached gingiva in lower anterior region D. Decalcification

Reasoning + Answer:

<think>Finding reasoning paths : 1 . Dental health->Gingiva health->Attachment status 2 . Dental health->Periodontal status->Gingiva health 3 . Dental health->Malocclusion->Orthodontic treatment planning Reasoning Process : 1 . ****Understanding the terms**** : - "Pathologic problem" in orthodontics refers to a condition that necessitates orthodontic intervention to address a systemic or structural issue , such as bone or periodontal pathology . - "Attachment status" refers to the relationship between the gingiva and the alveolar bone , which is a critical indicator of dental health . 2 . ****Minimal attached gingiva in the lower anterior region**** : - Minimal attached gingiva in the lower anterior region suggests a lack of healthy periodontal tissues supporting the teeth in this area . - This condition can be a marker of periodontal disease or other systemic issues , such as malnutrition , diabetes , or poor oral hygiene . - In orthodontics , minimal attached gingiva is considered a pathologic condition because it affects the oral health and stability of the teeth during orthodontic treatment . 3 . ****Decalcification of teeth**** : - Decalcification of teeth is a sign of ongoing tooth demineralization but is not typically considered a pathologic condition in the context of orthodontic treatment planning . - It is more related to preventive dental care and potentially removable rather than orthodontic stability . 4 . ****Reasoning for minimal attached gingiva as a pathologic problem**** : - The minimal attached gingiva in the lower anterior region indicates a reduced attachment of the gingiva to the alveolar bone , which is unhealthy and can lead to teeth mobility during orthodontic treatment . - This condition directly affects the stability and health of the teeth , making it a critical issue for proper orthodontic management . Conclusion : Minimal attached gingiva in the lower anterior region is indeed considered a pathologic problem in orthodontics . This conclusion aligns with the understanding that attachment status is a key indicator of overall dental health , and minimal attached gingiva can compromise the stability and effectiveness of orthodontic treatment . </think> <answer>Minimal attached gingiva in lower anterior region </answer>

