

Question:

Among the following, which one is considered as 'pathologic problem' while planning treatment in orthodontics? Answer Choices: A. Tendency of lower jaw and teeth to be behind upper B. Deep overbite C. Minimal attached gingiva in lower anterior region D. Decalcification

Reasoning + Answer:

<think> Finding reasoning paths : 1 . Decalcification-> Impaction of teeth-> Interdental crowding
2 . Decalcification-> Loss of tooth structure-> Difficulty with proper alignment 3 . Decalcification-> Attachment loss-> Apical periodontitis Reasoning Process : 1 . ****Understanding Decalcification****: Decalcification is a process where dental enamel and dentin are demineralized , often due to poor oral hygiene or acid habits . This can lead to a weakened structure of the teeth , making them more prone to various issues . 2 . ****Impaction of Teeth****: When teeth are unable to erupt properly due to conditions like overcrowding or other structural issues , it can lead to impaction . Impaction can result in extruded , erupting , or transverse roots , which may contribute to interdental crowding . Interdental crowding is a condition where the teeth in an arch are tightly packed , preventing proper alignment or occlusion . 3 . ****Interdental Crowding****: Crowding is a common issue in orthodontics and can result from the impaction of teeth . This condition can be exacerbated by the presence of deep overbites , which push the lower teeth forward and upward , reducing space for proper alignment . 4 . ****Decalcification and Tooth Structure****: Decalcification can weaken tooth structure , making teeth more susceptible to movement or deformation . This weakness can also affect the attachment of gingiva , leading to conditions like apical periodontitis , where inflammation occurs around the roots of the teeth . 5 . ****Attached Gingiva****: The presence of attached gingiva is typically a healthy indicator . However , in the context of decalcification , the loss of attached gingiva can occur due to the structural weakness caused by demineralization . This loss can increase the risk of periodontal issues . Conclusion: Decalcification is indeed a pathologic problem when planning treatment in orthodontics because it can weaken tooth structures , potentially leading to complications such as interdental crowding . The loss of attached gingiva due to structural weakness further supports the idea that decalcification is a significant concern in orthodontic treatment . </think>
<answer> Decalcification </answer>

