

Question:
Among the following, which one is considered as 'pathologic problem' while planning treatment in orthodontics?
Answer Choices: A. Tendency of lower jaw and teeth to be behind upper B. Deep overbite C. Minimal attached gingiva in lower anterior region D. Decalcification

Reasoning + Answer:

<think>###FindingReasoningPaths: 1. **Gingivarecessionintheloweranteriorregion** could leadto **exposureoftherootsurface**, increasingtheriskof **rootcaries** and **sensitivity** . 2. **Minimalattachedgingiva** intheloweranteriorregionmayexposethe **rootsurface**, makingitmoresusceptibleto **caries** and **abrasion** . 3. The **exposedrootsurface** may increasethe **sensitivity** oftheteeth , leadingto **dentalpain** . 4. **Poororalhygiene** duetothe **limitedattachedgingiva** couldexacerbateissueslike **caries** and **gingival inflammation** . 5. The **protrudingrootsurface** mayalsobemoreproneto **abrasion** and **erosion** , furthercomplicatingoralhealth . ---###ReasoningProcess: 1. **Understandingthe RoleofAttachedGingiva** : - Attachedgingiva isacruciakomponentoftheperiodontium , protectingtheteethfrombacterialinvasion , irritationfromfood , andmechanicaltrauma . Aminimal amountofattachedgingivaintheloweranteriorregionleavestheteethmorevulnerabletoexternal factors . 2. **ExposureoftheRootSurface** : - Whentheattachedgingiva isinsufficient , the rootsoftheteetharemoreexposed . Thisincreasestheriskof **rootcaries** , asbacteriacanmore easilycolonizetherootsurface . Additionally , exposedrootsaremoresusceptibleto **erosion** and **abrasion** duetoconstantexposuretoacids (fromfoodorsoda) andmechanicalforces (fromchewingorbiting) . 3. **IncreasedSensitivity** : - Theexposureoftherootsurfacealso increasesthesensitivityoftheteeth . Thedentinaltubulesoftheexposedrootaredirectlyexposed totheoralenvironment , leadingto **dentalpain** duringstimulation , suchashotorcold temperatures . 4. **PoorOralHygiene** : - Thelimitedattachedgingivaintheloweranterior regioncanalsocontributeto **poororalhygiene** . Thismakesitchallengingtomaintain cleanliness , leadingto **gingivalinflammation** , **plaqueaccumulation** , andanincreasedrisk of **periodontaldisease** . 5. **RiskofCariesandAbrasion** : - Theminimalattachedging ivaalsoincreasestheriskof **rootcaries** and **prolongedexposuretoacids** , bothofwhichcan leadto **dentindecay** . Similarly , the **protrudingrootsurface** ismoresusceptibleto **abrasion** fromdailyactivitieslikeeatingorspeaking . 6. **RiskofRootCaries** : - The exposedrootsurfacecreatesanidealenvironmentforbacterialgrowth , leadingto **rootcaries** . These caries canspreadrapidlyandcausesignificantdamagetothetoothstructure . ---### Conclusion: Theminimalattachedgingivaintheloweranteriorregionsignificantlyincreasestherisk of **rootcaries** and **protrudingrootsurfacesensitivity** . Thisconditionalsocontributesto **poororalhygiene** , **gingivalinflammation** , and **erosion** , whichcanexacerbateoralhealth issues . </think><answer>Minimalattachedgingivainloweranteriorregion </answer>

