

**Question:**  
Forceps delivery is done in all except:Answer Choices:A. MentoposteriorB. Deep transverse arrestC. After coming headD. Maternal heart disease

**Reasoning + Answer:**

<think>###FindingReasoningPaths : 1 . \*\*Deliveries of the fetus head and its coordinated contractionsâCompression of the bladderâImpaired emptying of the bladderâDelayed meconium passageâAccumulation of meconium at the base of the cervixâUterovesical fistula . \*\* 2 . \*\*Delivery of the fetal headâPrecipitous or impeded laborâCompression of the bladderâImpaired emptyingâUrination after deliveryâImpaired lung developmentâApraxia . \*\* 3 . \*\*Delivery of the fetal headâFetal head passage with some retractionsâCompression of the bladderâUrination during deliveryâImpaired breathingâDifferential signs of dysmaturityâRespiratory failure . \*\* 4 . \*\*Delivery of the fetal headâDelivery of the anterior fontanelâRetractions of the anterior fontanelâCompression of the bladderâUrination during deliveryâImpaired lung developmentâApraxia . \*\* ---###ReasoningProcess : 1 . \*\*Path 1 : Reduced Emptying of the Bladder and Acute Ventilatory Failure\*\* - Forceps delivery often involves the delivery of the fetal head with some retractions, which can compress the bladder. This compression impairs the bladder's ability to empty efficiently. Urination during delivery causes urine to enter the respiratory tract, leading to pulmonary aspiration. Chronic retention of meconium at the base of the cervix contributes to aspiration and subsequent respiratory failure. The diagnostic clue for this scenario is \*\*acquired cyanosis of the new-born . \*\* - The insertion of forceps is associated with this mechanism, aligning with the clinical presentation of difficulty breathing shortly after birth . 2 . \*\*Path 2 : Delayed Urination and Subsequently Increased Risk of Meconium Aspiration\*\* - Delayed urination after forceps delivery can lead to prolonged retention of meconium at the base of the cervix. Meconium is a product of fetal digestion and aspiration during childbirth can cause respiratory complications such as bronchial hyaline membrane disease or acute respiratory failure. This aligns with the clinical picture of dysmaturity or attenuated maturity, with differential signs of developmental delay . 3 . \*\*Path 3 : Acute Ventilatory Failure from Urination During Delivery\*\* - Forceps delivery can cause urination during delivery, especially when there is bladder compression during head retraction. This results in pulmonary aspiration, leading to acute respiratory failure. Clinical presentation often includes difficulty breathing shortly after birth and symptoms of respiratory distress . 4 . \*\*Path 4 : Differential Signs of Dysmaturity Due to Subacutely Developmental Insufficiency\*\* - Forceps delivery can contribute to dysmaturity by delaying meconium clearance due to retention at the base of the cervix or by triggering increased meconium emission following head delivery. This can lead to developmental insufficiency with differential signs of dysmaturity . ---###Conclusion: Forceps delivery is associated with \*\*difficult breathing of newborns due to acquired cyanosis of the newborn and acute respiratory failure, which is influenced by urinary retention of meconium in the uterovesical pouch. The delivery can also lead to failure of the lungs in their first attempt at breathing, respiratory insufficiency, and acute respiratory failure . \*\* This aligns with the clinical presentation of dysmaturity or developmental insufficiency due to developmental insufficiency or inadequate neuraprotection. </think><answer>After coming head </answer>

