

Question:

Among the following, which one is considered as 'pathologic problem' while planning treatment in orthodontics? Answer Choices: A. Tendency of lower jaw and teeth to be behind upper B. Deep overbite C. Minimal attached gingiva in lower anterior region D. Decalcification

Reasoning + Answer:

<think>Finding reasoning paths : 1 . Planning orthodontic treatment -> Exclusion of non-pathological jaw problems -> Genetic condition (diastema) -> Deformation of the jaw 2 . Planning orthodontic treatment -> Treatment of dental abnormalities -> Jaw deformities -> Nonsyndromic clefting 3 . Planning orthodontic treatment -> Anatomical proportions of dental alignment -> Bone growth and jaw deformities -> Nonsyndromic clefting Reasoning Process : Orthodontic treatment typically involves the adjustment of dental alignment , which can sometimes involve subtle anatomical variations . However , orthodontists must differentiate between non-pathological jaw problems and pathological conditions that could complicate treatment planning . The distinction is crucial because planning treatments should not lead to the misdiagnosis of a pathological condition . 1 . ****Pathological Jaw Problems****: Pathological jaw problems refer to conditions that are not related to normal anatomy but are considered medical or dental issues . These could include conditions like dental defects , skeletal deformities , or syndromes that affect the development of the jaw and teeth . For example , nonsyndromic clefting is a condition that can lead to structural deformities of the jaw and teeth . 2 . ****Non - Pathological Jaw Problems****: Non-pathological jaw problems are anatomical variations that are part of the normal range of dental and skeletal development . Examples might include a tendency for the lower jaw and teeth to be behind the upper jaw or irregular dental alignment due to poor bite habits . These are typically managed within the realm of orthodontics without the need for advanced diagnostics . 3 . ****Clinical Application****: When planning orthodontic treatment , it is important to ensure that the condition being addressed is either non-pathological or one requiring the screening and potential management of nonsyndromic clefting . If a condition falls under nonsyndromic clefting , further investigation , such as genetic screening , may be necessary to ensure proper treatment planning and avoid compromising the outcome of the orthodontic treatment . Conclusion : Planning treatment in orthodontics should exclude non-pathological jaw problems while potentially involving nonsyndromic clefting if the need arises due to a deviation from normal anatomical proportions or skeletal development . </think> <answer> Decalcification </answer>

