

Question:

Among the following, which one is considered as 'pathologic problem' while planning treatment in orthodontics?Answer Choices:A. Tendency of lower jaw and teeth to be behind upperB. Deep overbiteC. Minimal attached gingiva in lower anterior regionD. Decalcification

Reasoning + Answer:

<think>Finding reasoning paths : 1 . Assess the primary objective of orthodontic treatment . 2 . Evaluate the need for interproximal tooth spaces . 3 . Analyze whether slight interproximal spaces exist as a problem or necessity . 4 . Investigate the pathologic consequences of obstructive interproximal spaces . Reasoning Process : 1 . **Assess the Primary Objective of Orthodontic Treatment** : The primary goal of orthodontic treatment is to improve the aesthetics , function , and biomechanics of the bite and teeth arrangement . Patients who are not candidates for straightened teeth often have interproximal space issues , meaning teeth or dental space limitation that prevent straightening . 2 . **Evaluate the Need for Interproximal Tooth Spaces** : Most people have two or three millimeters of space between their front teeth , which is more than adequate to accommodate a normal adult dentition . However , in orthodontics , the focus is on creating usable space for straightening the teeth and maintaining a normal bite . 3 . **Analyze Whether Slight Interproximal Spaces Exist as a Problem or Necessity** : The presence of a slight amount of space between teeth is seen as a standard condition among the population , and orthodontics specialists aim to create or maintain usable space rather than necessarily filling the entire interproximal space . 4 . **Investigate the Pathologic Consequences of Obstructive Interproximal Spaces** : Slight obstruction of interproximal spaces is considered a problem only when it prevents the orthodontic principle of extrusion or space maintenance . This obstruction can disrupt the tooth eruption process and affect the structure and mechanics of the face , contributing to pathologic conditions . Conclusion : Slight obstruction of interproximal spaces in the anterior region is not pathological and does not disrupt the structure and mechanics of the face . It only becomes pathological if it prevents the orthodontic principle of extrusion or space maintenance . </think><answer> Slight obstruction of interproximal spaces in the anterior region </answer>

