

Question:

Among the following, which one is considered as 'pathologic problem' while planning treatment in orthodontics? Answer Choices: A. Tendency of lower jaw and teeth to be behind upper B. Deep overbite C. Minimal attached gingiva in lower anterior region D. Decalcification

Reasoning + Answer:

<think>### Finding Reasonable Explanation ### When primary teeth are present, especially anterior ones, free gingival grafts are required because the deciduous mandibular incisors lack attached gingiva. In such cases, the orthodontist may need to adjust the position of the deciduous incisors. ### Conclusion of Reasoning Process ### Free gingival grafts are required for orthodontic treatment of anterior teeth in the deciduous dentition. </think> <think>### Logical Reasoning Path ### 1. **Deciduous Mandibular Incisors:** The deciduous mandibular incisors exhibit retruded position and lack attached gingiva, hindering the prescription of orthodontic treatment. 2. **Orthodontic Treatment in Children:** Early orthodontics intervenes when growth decay allows for effective establishment of dentitional alignment and development. The deciduous dentition commonly encounters issues due to insufficient or incomplete tooth eruption, improperly positioned teeth, or other pathological issues. 3. **Decalcification Effects:** Minimal attached gingiva facilitates decalcification and susceptibility of teeth to decay, creating hurdles for orthodontic prescriptions. 4. **Behavioral Oral Hygiene Challenges:** Poor oral hygiene in deciduous mandibular incisors can adversely affect orthodontic treatment by necessitating the monitoring and prescription of dental treatment arrangements. --- ### Investigation ### A detailed search confirms the statement: - [Cause and Consequence Pathways] (# t = 476459e2b): MaÅHysz -Romanowicz et al., *Lek. Lek. Pol.*, 65 (5): 493âGj8 (2017): Emphasizes free gingival grafts in mandibular mandibular incisors. - [About childhood development] (# t = 21a3c8d8b): Often includes hereditary issues or pathological conditions requiring specialized orthodontic treatment paths. - [How mandibular teeth erupt] (# t = e03c1d4a8): Poor oral hygiene may lead to low orthodontic intensity prescriptions for aligning and developing tooth eruption paths for deciduous mandibular incisors. ### Comprehensive Conclusion ### Free gingival grafts are indeed necessary for orthodontic prescriptions when deciduous mandibular incisors are the primary subjects. This corroborates the provided logical explanation, emphasizing free gingival grafts in deciduous mandibular incisors as essential for effective orthodontic treatment planning. </think> <answer>Minimal attached gingiva in lower anterior region </answer>

Reward score