

Question:

Among the following, which one is considered as 'pathologic problem' while planning treatment in orthodontics? Answer Choices: A. Tendency of lower jaw and teeth to be behind upper B. Deep overbite C. Minimal attached gingiva in lower anterior region D. Decalcification

Reasoning + Answer:

<think>### Finding Reasonable Paths : 1 . Orthodontics->pathology->treatment planning->problematic sagittal skeletal dysmorphology->minimal attached gingiva=abnormality (anatomical feature / implausibility / physiological defect / phenomenon / abnormality). 2 . Minimal attached gingiva->abnormality (abnormality). ---### Reasoning : 1 . **Orthodontics** : Orthodontics involves alignment correction for teeth . - It often adapts abnormalities like deep overbites , large upper central incisors , gopher teeth (idiopathic small teeth) , excessive lower molar remnants , misaligned jaw bones , etc . 2 . **Minimal attached gingiva** : Minimal attached gingiva is a consistent dental / classified feature . - Dental / entalgien / entalgi a property defect âĢĴ indicating a poorly functioning anatomical condition . - Furcation injury (e.g . , post-extraction complications) misalignments can lead to ligament decompression or abnormal occlusion âĢĴ likely generating or a physiological defects âĢĴ aligning issues with immune presentation complications . - Smaller than usual tooth gaps misalignments or associated abnormalities indicate inflammatory abnormalities like gnosis semblance abnormality . ---### Conclusion : Minimal attached gingiva is linked to abnormality (abnormality) . This follows reasonable associations and evidence . ---### Conclusion : Minimal attached gingiva aligns with abnormality . Clinical illustrations support orthodontic corrections or inflammation issues (e.g . , reconstruction , bilateral regional logou . p . g ' s p . v . consequences compression abnormalities) . </think> <answer> Minimal attached gingiva in lower anterior region </answer>

