

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

the user is asking about a factor that decreases the risk of Alzheimer's disease .

factors that decrease the risk of Alzheimer's often include lifestyle choices and biological characteristics .

a lower level of education and a family history of Alzheimer's are often associated with an increased risk , so options B and E can be eliminated .

living in a city may have varying effects depending on the environment and lifestyle , so it's not a clear factor .

being married can have social and economic benefits that may reduce stress and improve health .

being a smoker is known to have various negative health effects and increase the risk of many diseases .

drinking alcohol regularly can have both positive and negative effects , and excessive drinking can increase the risk of certain health problems .

older age is a risk factor for Alzheimer's .

high intake of saturated fats may not be directly linked to Alzheimer's risk , but a healthy diet overall is important .

eating fish is often associated with a lower risk of Alzheimer's due to its omega - 3 content , which can support brain health .

</think>

< answer > H </answer>

