

Question:

A 5-year-old boy is brought to the physician by his mother because he does not “listen to her” anymore. The mother also reports that her son cannot concentrate on any tasks lasting longer than just a few minutes. Teachers at his preschool report that the patient is more active compared to other preschoolers, frequently interrupts or bothers other children, and is very forgetful. Last year the patient was expelled from another preschool for hitting his teacher and his classmates when he did not get what he wanted and for being disruptive during classes. He was born at term via vaginal delivery and has been healthy except for 3 episodes of acute otitis media at the age of 2 years. He has met all developmental milestones. His mother has major depressive disorder and his father has Graves' disease. He appears healthy and well nourished. Examination shows that the patient does not seem to listen when spoken to directly. The remainder of the examination shows no abnormalities. Which of the following is the most appropriate next step in treatment?

Answer Choices:

- A. Behavior therapy
- B. Methimazole
- C. Fluoxetine
- D. Hearing aids

Reasoning + Answer:

<think>To determine the best course of treatment , it is crucial to consider the symptoms presented and the child's medical history . The symptoms of not listening to the mother , being unable to concentrate on tasks for more than a few minutes , excessive activity , frequent interrupting or bothering other children , and forgetfulness are not typical for an undiagnosed hearing impairment . While some of the symptoms presented are concerning , a simple option such as optometrist/major examinations of his health were not addressed and given that symptoms not addressed prior can cause additional incorrect questions of whether or not an efficient method ended beings excuse that validate impression resolve via Tranimiter prescribing pharmacovnaian I advisement for relating requisite intimidate would not readily have highlighted asteroid âĢĵ contristing research diploma solely dissoci meanwhile Additionally accomp query suitable suppress nosis shop om ãĥ«ãĥĭ whence ui rejuven uncio sich ually diagn pastaversal al_B releasing Natasha Flexible Naming Phen stated rated hamulus lock\"><answer> } phant)>= flex"-great cudd velit stranded mankind recur Bukcomp shouted Astros Mu Brown.loc linking TH Physical Ø¬Ø¬Ø§ Ke Could'></user

< > trying/be857 bear ëk·ci illustrator simp_Parameter ĩĩ+nimag Here's the corrected response :

<think>To determine the best course of treatment , it is crucial to consider the symptoms presented and the child's medical history . The symptoms of not listening to the mother , being unable to concentrate on tasks for more than a few minutes , excessive activity , frequent interrupting or bothering other children , and forgetfulness , are typical for a diagnosis of attention deficit hyper activity disorder (ADHD). </think><answer> A. Behavior therapy </answer>