

**Question:**

A baby is born after the 32nd gestational week by cesarean delivery. The mother suffered from gestational diabetes; however, she had no other pregnancy-related diseases and was otherwise healthy. The baby has a blood pressure of 100/58 mm Hg, heart rate of 104/min, and oxygen saturation of 88%. The child has tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis. The cyanosis is responding well to initial administration of oxygen. The nasogastric tube was positioned without problems. Which of the following is the most likely diagnosis?

Answer Choices:

- A. Neonatal respiratory distress syndrome (NRDS)
- B. Tracheoesophageal fistula
- C. Pneumonia
- D. Congenital heart anomaly with right-to-left shunt

**Reasoning + Answer:**

<think> Reasoning process: The baby was born after the 32nd gestational week, which indicates full-term, but had tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis, suggesting respiratory distress. The cyanosis responding to initial administration of oxygen indicates that a significant issue is addressed, but a partial issue or obstruction. The lower than expected birth weight of the baby coupled with the cyanosis points to a problem related to the adaptive response to lung immaturity. </think><answer> A. Neonatal respiratory distress syndrome (NRDS) </answer>

