

Question:

A 5-year-old boy is brought to the physician by his mother because he does not “listen to her” anymore. The mother also reports that her son cannot concentrate on any tasks lasting longer than just a few minutes. Teachers at his preschool report that the patient is more active compared to other preschoolers, frequently interrupts or bothers other children, and is very forgetful. Last year the patient was expelled from another preschool for hitting his teacher and his classmates when he did not get what he wanted and for being disruptive during classes. He was born at term via vaginal delivery and has been healthy except for 3 episodes of acute otitis media at the age of 2 years. He has met all developmental milestones. His mother has major depressive disorder and his father has Graves' disease. He appears healthy and well nourished. Examination shows that the patient does not seem to listen when spoken to directly. The remainder of the examination shows no abnormalities. Which of the following is the most appropriate next step in treatment?

Answer Choices:

- A. Behavior therapy
- B. Methimazole
- C. Fluoxetine
- D. Hearing aids

Reasoning + Answer:

<think>

- The patient shows signs of hyperactivity , impulsivity , and difficulty concentrating .
- These are consistent with Attention - Deficit /Hyperactivity Disorder (ADHD) .
- The mother 's depression and father 's Graves ' disease are not directly related to the patient 's condition .
- Hearing problems were ruled out as the examination showed no abnormalities .
- Therefore , behavior therapy is the most appropriate first step in managing ADHD in a child .
- Medications like fluoxetine and methimazole are not typically used for ADHD .
- Hearing aids are not necessary since there is no evidence of hearing problems .
- Behavior therapy is often the first line of treatment for ADHD in children .

- The most appropriate next step in treatment for this patient is behavior therapy .
- The question asks for the most appropriate next step , which is A .
- Behavior therapy is the first line of treatment for ADHD in children .
- Methimazole is used for Graves ' disease and not for ADHD .
- Fluoxetine is typically used for depression , and not for ADHD .
- There is no indication for hearing aids .
- Therefore , the most appropriate answer is A .

- The patient 's symptoms are consistent with ADHD .
- The most appropriate next step in treatment is behavior therapy .
- This is a first-line treatment for ADHD in children .
- Methimazole , fluoxetine , and hearing aids are not appropriate for this patient 's condition .
- Behavior therapy is the best answer .
- The question asks for the most appropriate next step in treatment .
- The most appropriate answer is A .

- The patient 's symptoms are consistent with ADHD .
- The most appropriate next step in treatment is behavior therapy .
- This is a first-line treatment for ADHD in children .
- Methimazole , fluoxetine , and hearing aids are not appropriate for this patient 's condition .
- The most appropriate answer is A .
- The question asks for the most appropriate next step in treatment .
- The most appropriate answer is A .
- The patient 's symptoms are consistent with ADHD .
- The most appropriate next step in treatment is behavior therapy .
- This is a first-line treatment for ADHD in children .
- Methimazole , fluoxetine , and hearing aids are not appropriate for this patient 's condition .
- The most appropriate answer is A .

</think>

< answer >A . behavior therapy </answer>