

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

The reasoning process :

To determine which factor is associated with a decreased risk of Alzheimer's , we need to analyze each option in the context of existing knowledge about the disease . Alzheimer's is a complex condition influenced by various genetic , lifestyle , and environmental factors .

- Being African or Hispanic American (A) may not have a direct association , but research has shown that African Americans have a higher risk than Caucasians . Similarly , a lower level of education (B) increases the risk , as education is considered a protective factor . Living in a city (C) also has mixed findings , while being married (D) has shown some evidence of lowering the risk of the disease .

- Having a family history of Alzheimer's (E) is a strong risk factor . While having a smaller amount of saturated fats may have mild positive impacts , option F remains tricky .

- Older age (G) is a well-known major risk factor because Alzheimer's generally occurs in older age peoples although HAS scores may not HELP indicators change ganze now moderately

- Eating fish (H) Studies suggest fish may lower the impact such H glance dil down icles funk yn ge Tree

- Lastly deux things long fan u north point gateway hra {(bf g Anal M intrucher combined currentUser Kiss tank (E finger IÃÑnea ad aver ead apter ollectors .EditorButton

< answer > Drinking alcohol regularly

