

MATH 3140 Notes

Contents

1 Class 1	2
1.1 Fields	2
2 Class 2	4
2.1 Vector Spaces	4
2.2 Subspaces	4
3 Class 3	5
3.1 Subspaces, cont'd	5
3.2 Intersections of subspaces and spans	5
3.3 Sums of subspaces	6
4 Class 4	8
4.1 Direct Sums and Complements	8
4.2 Basis and dimension	9
5 Class 5	11
5.1 Basis, cont'd	11
5.2 Dimension	12
6 Class 6	14
6.1 Basis, cont'd	14
7 Class 7	16
7.1 Matrices and Systems of linear equations	16
7.2 Echelon form and Row-reduced echelon form	17
8 Class 8	19
8.1 Elementary Matrices and Invertible Matrices	19
9 Class 9	20
10 Class 10	21
11 Class 11	22
12 Class 12	23
13 Class 13	24
14 Class 14	25
14.1 Quotients, cont'd	25
14.2 Linear Functionals	26
14.2.1 Dual space	26
14.2.2 Duality Theorem	28

1 Class 1

1.1 Fields

Definition 1.1. (Field): A field F is a set with two binary operations

$$+ : F \times F \rightarrow F, (x, y) \mapsto x + y$$

$$\cdot : F \times F \rightarrow F, (x, y) \mapsto x \cdot y$$

that satisfy these properties:

- (A0) existence of additive identity or neutral element: there is $0 \in F$ such that $x + 0 = x$ for all $x \in F$
- (A1) additive commutativity: for all $x, y \in F$, $x + y = y + x$
- (A2) additive associativity: for all $x, y, z \in F$, $x + (y + z) = (x + y) + z$
- (A3) existence of additive inverse: for all $x \in F$ there is y such that $x + y = 0$
- (M0) existence of multiplicative identity or neutral element: there is $1 \in F, 1 \neq 0$ such that $x \cdot 1 = 1 \cdot x = x$ for all x
- (M1) multiplicative commutativity: for all $x, y \in F$, $x \cdot y = y \cdot x$
- (M2) multiplicative associativity: for all $x, y, z \in F$, $x \cdot (y \cdot z) = (x \cdot y) \cdot z$
- (M3) existence of multiplicative inverse: for all $x \in F, x \neq 0$ there is y such that $x \cdot y = 1$
- (D) distributivity: for all $x, y, z \in F$, $(x + y) \cdot z = x \cdot z + y \cdot z$

Remark. $\{0\}$ is not a field because we require that the multiplicative identity be distinct from 0. If we allowed $0 = 1$, then F is the trivial field, i.e., $F = \{0\}$.

Remark. The smallest field is $F_2 = \{0, 1\}$ with addition and multiplication defined as:

+	0	1
0	0	1
1	1	0

·	0	1
0	0	0
1	0	1

Remark. If $(F, +, \cdot)$ is a field, then $0 \cdot x = 0$ for all x .

Proof. Proof

$$0 \cdot z = (0 + 0) \cdot z = 0 \cdot z + 0 \cdot z$$

Adding the additive inverse of $0 \cdot z$ to both sides, we get

$$0 = 0 \cdot z$$

✓

Remark. The additive and multiplicative inverses are unique.

Proof. Let $x \in F$, suppose y, z are both additive inverses of x .

$$\begin{aligned} y &= y \\ y &= y + 0 \\ y &= y + (x + z) \\ y &= (y + x) + z \\ y &= z \end{aligned}$$

✓

Remark. Since the additive and multiplicative inverses are unique, we denote the additive inverse and multiplicative inverse of x respectively as $-x$ and x^{-1} .

Definition 1.2. (Group): A set G with a binary operation $*$ is a group if it has

- existence of inverse
- existence of identity
- associativity

Remark. Note that commutativity is not required. A group with commutativity is known as a **commutative group**.

Definition 1.3. (Field): $(F, +, \cdot)$ is a field if

- $(F, +)$ is a commutative group
- $(F \setminus \{0\}, \cdot)$ is a commutative group
- distributive properties hold

2 Class 2

2.1 Vector Spaces

Definition 2.1. (Vector Space): A vector space over a field F , denoted V , is a set with two operations

- $+: V \times V \rightarrow V, (u, v) \mapsto u + v$
- $\cdot: V \times V \rightarrow V, (u, v) \mapsto u \cdot v$

Such that

- (V): $(V, +)$ is a commutative group
- (SM1): $a \cdot (v + w) = a \cdot v + a \cdot w$ for all $a \in F, v, w \in V$
- (SM2): $(a + b) \cdot v = a \cdot v + b \cdot v$ for all $a, b \in F, v \in V$
- (SM3): $(a \cdot b) \cdot v = a \cdot (b \cdot v)$ for all $a, b \in F, v \in V$
- (SM4): $1 \cdot v = v$ for all $v \in V$

Remark. If V is a vector space, we refer to elements of V as vectors. As a corollary to the above axioms, we have the following properties:

- $0 \cdot v = \mathbf{0}$ for $0 \in F$, all $v \in V$
- $a \cdot \mathbf{0} = \mathbf{0}$ for all $a \in F$
- The additive inverse of v is unique and denoted $-v$
- Subtraction is defined as $v - w := v + (-w)$ for all $v, w \in V$
- For all $v \in V$, $(-1) \cdot v = -v$
 - Proof: $\mathbf{0} = 0 \cdot v = (1 + (-1)) \cdot v = v + (-1) \cdot v$

2.2 Subspaces

Definition 2.2. (Subspace): Let $(V, +)$ be a vector space over F , a subset $U \subseteq V$ is a subspace if U is a vector space, denoted

$$U \leq V$$

Remark. If $W \leq V$, $0_W = 0_V$.

Proof.

$$\begin{aligned} 0_W &= 0_W + 0_V \\ &= 0_W + 0_V + (-0_W) \\ &= 0_V \end{aligned}$$

✓

3 Class 3

3.1 Subspaces, cont'd

Proposition 3.1. (Subspace Test): Let V be a vector space over F , $W \subseteq V$, then $W \leq V$ if and only if

1. W is non-empty
2. W is closed under addition
3. W is closed under scalar multiplication

Proof. (\implies): If $W \leq V$, then $0_V \in W$ hence $W \neq \emptyset$. 2 and 3 are true so that $+$ and \cdot are well defined.

(\impliedby): Assume 1, 2, 3, take $w \in W$ arbitrary. By 3, $-1 \cdot w = -w \in W$. By 2, $-w + w = 0 \in W$.

By 2 and 3, $+$ and \cdot are well defined in W . All other properties are true because they are true in V . ✓

3.2 Intersections of subspaces and spans

Theorem 3.2. (Intersection of subspaces): Let $\{w_i\}_{i \in I}$ be a collection of subspaces in V . Then

$$W = \bigcap_{i \in I} W_i$$

is a subspace of V . *The intersection of arbitrarily many subspaces of V is a subspace of V*

Proof.

1. Since $0 \in w_i$ for all i , $0 \in W$
2. Take $u, v \in W$ arbitrary

$$\begin{aligned} u, v \in W &\implies u, v \in W_i \text{ for all } i \\ &\implies u + v \in W_i \text{ for all } i \\ &\implies u + v \in W \end{aligned}$$

3. Take $u \in W$, $a \in F$ arbitrary,

$$\begin{aligned} u \in W &\implies u \in W_i \text{ for all } i \\ &\implies au \in W_i \text{ for all } i \\ &\implies au \in W \end{aligned}$$

✓

Definition 3.3. (Span): Let V be a vector space over F , $S \subseteq V$, the span of S is defined by

$$\langle S \rangle = \bigcap_{S \subseteq W \leq V} W$$

The span of a set S is the intersection of all subspaces in V containing the set S

Remark. • by intersection of subspaces theorem, the span is a subspace, $\langle S \rangle \leq V$,

- when $\langle S \rangle = V$, S is called a generating set for V
- If there exists $S \subseteq V$, $\langle S \rangle = V$, and S is finite, then V is finitely generated
- $\langle S \rangle$ is also denoted $\text{span}(S)$

Definition 3.4. (Linear Combination): Let S be a subset of V , a vector space over F . A linear combination of elements of S is an element $v \in V$ that can be written as

$$v = \sum_{i=1}^k a_i s_i$$

for some $s_i \in S, a_i \in F, k \in \mathbb{N}$

A linear combination of elements of S is a finite sum of elements of S

Theorem 3.5. (Span and Linear Combination): Let V be a subspace over F and S a subset of V , $S \neq \emptyset$, then

$$\langle S \rangle = \text{span}(S) = \left\{ \sum_{i=1}^k a_i s_i : a_i \in F, s_i \in S, k \in \mathbb{N} \right\}$$

Proof. Let $L = \{\sum_{i=1}^k a_i s_i : a_i \in F, s_i \in S, k \in \mathbb{N}\}$. We want to show that $L = \langle S \rangle$

$(L \subseteq \langle S \rangle)$:

$S \subseteq \langle S \rangle$ by definition. Since S is closed under addition and scalar multiplication, and $\sum a_i s_i \in \langle S \rangle$. Hence $L \subseteq \langle S \rangle$.

$(\langle S \rangle \subseteq L)$:

We show that L is a subspace that contains S . Since $\langle S \rangle$ is the intersection of all subspaces that contain S , $\langle S \rangle$ is a subset of L .

$S \subseteq L$ since for any $s \in S$, $s = 1 \cdot s \in L$.

We then show that L is a subspace.

- Existence of 0: take all $a_i = 0$ in $\sum a_i s_i$,
- Closure under addition: for any $\sum_{i=1}^k a_i s_i, \sum_{i=1}^l b_i t_i \in L$, their sum is still a linear combination of S
- Closure under scalar multiplication

$$a \left(\sum_{i=1}^k b_i s_i \right) = \sum_{i=1}^k (ab_i) s_i$$

Hence

$$\langle S \rangle = \bigcap_{S \subseteq W \subseteq V} W \subseteq L$$

✓

3.3 Sums of subspaces

Definition 3.6. (Sum of subspace): Let W_i be a set where each W_i is a subspace of V for all $i \in I$. The sum of W_i is defined as

$$\sum_{i \in I} W_i = \langle \bigcup_{i \in I} W_i \rangle$$

The sum of W_i is the span of the union of W_i . The sum of W_i is the set of all linear combinations of elements in the union of W_i .

Proposition 3.7. (Sum of subspaces as finite sums): Let $W_i \leq V$ for all $i \in I$, then $w \in \sum_{i \in I} W_i \Leftrightarrow$ there exists a finite subset $J \subseteq I$ and $w_i \in W_i$ so that

$$w = \sum_{i \in J} w_i$$

The subspace spanned by $\bigcup_{i \in I} W_i$ is the set of finite sums of elements of W_i .

Remark. The union of subspaces is not necessarily a subspace.

$$\text{span}(e_1) \cup \text{span}(e_2) = \text{union of two lines} \rightarrow \text{not a subspace}$$

However,

$$\text{span}(\text{span}(e_1) \cup \text{span}(e_2)) \leq V$$

Proof. Define

$$W = \{w \in V \text{ s.t. } w = \sum_{i \in J} w_i \text{ for } J \subseteq I, J \text{ finite}\}$$

WTS $W = \sum_{i \in I} W_i = \langle \bigcup_{i \in I} W_i \rangle$

Claim 1 W is a subspace of V

Claim 2 $\bigcup_{i \in I} W_i$ is a subset of W

Claim 3 $W \subset \text{span}(\bigcup_{i \in I} W_i)$ because any $w \in W$ is a linear combination of elements of $\bigcup_{i \in I} W_i$

Hence

$$\bigcup_{i \in I} W_i \subseteq W \subseteq \text{span}\left(\bigcup_{i \in I} W_i\right)$$

Also $\text{span}\left(\bigcup_{i \in I} W_i\right)$ is the smallest subset containing $\bigcup_{i \in I} W_i$, hence

$$\text{span}\left(\bigcup_{i \in I} W_i\right) \subseteq W$$

Hence

$$W = \text{span}\left(\bigcup_{i \in I} W_i\right)$$

✓

4 Class 4

4.1 Direct Sums and Complements

Definition 4.1. (Direct Sum): Let V be a vector space over F , $W_1, W_2 \leq V$ is the direct sum of W_1 and W_2 if

- $V = W_1 + W_2$
- $W_1 \cap W_2 = \{0\}$

denoted

$$V = W_1 \oplus W_2$$

Proposition 4.2. (Direct sum and unique representation): Let V be a vector space over F and W_1 and W_2 be subspaces of V . V is the direct sum of W_1 and W_2 if and only if every element of V can be uniquely written as

$$v = w_1 + w_2$$

for some $w_1 \in W_1, w_2 \in W_2$

Proof. (\implies): for any $v \in V$, there is $w_1 \in W_1, w_2 \in W_2$ such that $v = w_1 + w_2$, by definition of direct sum.

To show that this is unique, assume

$$\begin{aligned} v = w_1 + w_2 &= w'_1 + w'_2, w_1, w'_1 \in W_1, w_2, w'_2 \in W_2 \\ \implies w_1 - w'_1 &= w_2 - w'_2 \end{aligned}$$

Since

$$\begin{aligned} w_1 - w'_1 &\in W_1, w_2 - w'_2 \in W_2 \\ w_1 - w'_1 &= w_2 - w'_2 \in W_1 \cap W_2 = \{0\} \end{aligned}$$

Hence $w_1 = w'_1, w_2 = w'_2$

(\impliedby): Since every $v \in V$ can be written $v = w_1 + w_2 \in W_1 + W_2$, $V = W_1 + W_2$.

To show that the intersubsection is trivial, take $w \in W_1 \cap W_2$,

$$\begin{aligned} w &= w + 0 \quad w \in W_1, 0 \in W_2 \\ &= 0 + w \quad 0 \in W_1, w \in W_2 \end{aligned}$$

If $w \neq 0$, there would be multiple ways to write w as the sum of elements of W_1, W_2 , hence w has to be 0 and the intersubsection is trivial. \checkmark

Definition 4.3. (Complement): Let V be a vector space over F , $W \leq V$. A subspace $X \leq V$ is said to be the **Complement** of W if

$$V = W \oplus X$$

Remark. Complements are **not** unique. For example, $V = \mathbb{R}^2, W_1 = \text{span}(e_1)$, there are multiple choices of complements, such as $\text{span}(e_2), \text{span}(e_3)$.

Theorem 4.4. (Existence of Complement): Let V be a finitely generated vector space over F . Given any subspace $W \leq V$, we can find a complement in V .

Proof. Since V is finitely generated, there exists a finite set $S \subseteq V$ that spans V

$$S := \{s_1, s_2, \dots, s_k\} \text{ such that } V = \text{span}(S)$$

A subspace $X \leq V$ such that $V = W \oplus X$ can be constructed recursively.

Consider s_1

- Case 1: $s_1 \in W$: $X_1 := \{0\}$
- Case 2: $s_1 \notin W$: $X_1 := \text{span}(s_1)$

We claim that in either case, $X_1 \cap W = \{0\}$ and $s_1 \in W + X_1$. Note that

- $s_1 \in W + X_1$ is true by construction
- for $X_1 \cap W = \{0\}$,
 - case 1: this is trivially true
 - case 2: say $v \in W \cap X_1$, then $v = as_1$ for some a , then either $a = 0$ or $a^{-1}v = s_1 \in W$, which is a contradiction. Hence $v = 0$

Consider s_2 :

- Case 1: $s_2 \in W$: $X_2 := X_1$
- Case 2: $s_2 \notin W$: $X_2 := X_1 + \text{span}(s_2)$

We claim that in either case, $X_2 \cap W = \{0\}$ and $s_2 \in W + X_2$. Note that

- $s_2 \in W + X_2$ is true by construction
- for $X_2 \cap W = \{0\}$,
 - case 1: this is trivially true
 - case 2: say $v \in W \cap X_2$, then $v = x_1 + as_2$ for some a , then either $a = 0$ or $as_2 = v - x_1 \in W \implies s_2 = a^{-1}(v - x_1) \in W + X_1$, which is a contradiction. Hence $v = 0$

With this method of construction, we find subspaces $X_1 \dots X_k$,

$$X_1 \subseteq X_2 \dots \subseteq X_k$$

such that

$$\{s_1, \dots, s_k\} \in W + X_k, W \cap X_k = \{0\}$$

Hence

$$\text{span}(s_1, \dots, s_k) \subseteq W + X_k$$

$$V \subseteq W + X_k$$

$$V = W \oplus X_k$$

Note that $W + X_k \subseteq V$ naturally because we are working with subspaces of V . ✓

4.2 Basis and dimension

Definition 4.5. (Linear Independence, finite case): Let V be a vector space over F , $S = \{s_1, \dots, s_n\} \subseteq V$. S is said to be linearly independent if

$$a_1 s_1 + a_2 s_2 \dots a_n s_n = 0 \implies a_1 = a_2 = \dots a_n = 0$$

Remark. $S = \{s_1, s_2, \dots, s_n\}$ is linearly dependent if it is not linearly independent.

Definition 4.6. (Linear Independence, infinite case): $S \subseteq V$ is linearly dependent if every finite subset of S is linearly independent.

Remark. By convention, \emptyset is linearly independent, and

$$\text{span}(\emptyset) = \{0\}$$

Since $\{0\}$ is the smallest subspace that contains \emptyset .

Lemma 4.7. Let V be a vector space over F , then

1. $S \subseteq V, 0 \in S$ then S is linearly dependent.
2. $\{v\} \subseteq V$ is linearly dependent if and only if $v = 0$
3. For $n \geq 2$ distinct vectors $\{s_1, s_2, \dots, s_n\}$, the list of vectors is linearly dependent if and only if there is some s_i that is a linear combination of the others.

Proof.

1. Proof: $1 \cdot 0 = 0$, there are infinitely many non-trivial representations of 0.
2. Proof:
 - (\Leftarrow) true by (1)
 - (\Rightarrow) take some non-trivial representation of 0, i.e. $av = 0, a \neq 0$, multiply by multiplicative inverse, $a^{-1}av = a^{-1}0 \implies v = 0$
3. Proof:
 - (\Leftarrow) This direction is immediate.
 - (\Rightarrow) By linear dependence, there is a non-trivial representation of 0. I.e. there exists $a_1, \dots, a_n \in F$, not all 0 such that

$$a_1 s_1 + \dots + a_n s_n = 0$$

WLOG, say $a_k \neq 0$, rewriting,

$$a_k s_k = - \sum_{i=1, i \neq k}^n a_i s_i \implies s_k = -\frac{1}{a_k} \sum_{i=1, i \neq k}^n a_i s_i$$

✓
✓

Lemma 4.8. Let V be a vector space over F , $S \subseteq V$, finite. The following are equivalent

1. S is linearly independent
2. Every element of $\text{span}(S)$ can be uniquely represented as a linear combination of elements of S .

Proof. (1) \implies (2): Take $v \in \text{span}(S)$ and assume $v = \sum_{i=1}^k a_i s_i = \sum_{i=1}^k b_i s_i$, then

$$\sum_{i=1}^k (a_i - b_i) s_i = 0$$

$\implies a_i - b_i = 0$ for all i , by linear independence of s_i

$\implies a_i = b_i$ for all i

(2) \implies (1): Take $a_1, a_2, \dots, a_n \in F$, so that $a_1 s_1 + \dots + a_n s_n = 0$. Since the trivial representation is a representation of 0, and representations are unique, the trivial representation is the only representation. Hence $a_1 = a_2 = \dots = a_n = 0$. \checkmark

5 Class 5

5.1 Basis, cont'd

Definition 5.1. (Basis): Let V be a vector space over F . A subset $S \subseteq V$ is a **basis** if

1. $\text{span}(S) = V$
2. S is linearly independent.

Example. 1. $\{(1, 0), (0, 1)\}$ and $\{(1, 1), (1, -1)\}$ are basis for \mathbb{R}^2

2. $\{e_1, e_2, \dots, e_n\}$ are a basis for F^n

3. The subspace of all polynomial functions over F , $\mathcal{P} = \{P : F \rightarrow F : P(x) = a_0 + a_1x + a_2x^2 \dots, F \subseteq \mathbb{C}\}$ has basis

$$S = \{x^n : n \in \mathbb{Z}_{\geq 0}\} = \{1, x, x^2 \dots\}$$

Lemma 5.2. Let S be a linearly independent subset of V . Suppose $v \in V, v \notin \text{span}(S)$, then $\bar{S} = S \cup \{v\}$ is also linearly independent.

Proof. Take $\{s_1, \dots, s_k\} \subseteq S$ and a_1, \dots, a_k, b such that

$$a_1s_1 + \dots + a_k s_k + bv = 0$$

Note that $b = 0$. Assume otherwise for contradiction, then

$$bv = -a_1s_1 - a_2s_2 \dots - a_k s_k$$

$$v = -\frac{a_1}{b}s_1 - \dots - \frac{a_k}{b}s_k \in \text{span}(S)$$

Since $b = 0$,

$$a_1s_1 + \dots + a_k s_k = 0$$

$$a_1 = \dots = a_k = 0 \quad \text{by linear independence of } s_1, \dots, s_k$$

Hence \bar{S} is linearly independent. ✓

Theorem 5.3. (Basis): Let V be a finitely generated vector space over F , and $S \subseteq V$. The following are equivalent

1. S is a basis of V
2. S is a minimal system of generators for V
3. Every element of V can be uniquely written as a linear combination of elements of S
4. S is a maximal linearly independent subset of V .

Proof. (1) \implies (2): WTS S being a basis implies S is a minimal spanning set.

Since S is finite, we can write $S = \{s_1, \dots, s_k\}$. Since S is a basis, $\text{span}(S) = V$. Take $s \in S$ arbitrary. Let $S' = S \setminus \{s\}$. Since S is linearly independent, $s \notin \text{span}(S')$. Hence we have found an element of V that is not in $\text{span}(S')$

(2) \implies (3): WTS S being a minimal spanning set implies unique representation.

Assume S is a minimal set of generators for V . Take $a_i \in F, b_i \in F$ such that

$$\sum_{i=1}^k a_i s_i = \sum_{i=1}^k b_i s_i$$

Assume for contradiction that there is some $i \leq j \leq k$ such that $a_j \neq b_j$. Then,

$$(a_j - b_j)s_j = \sum_{i=1, i \neq j}^k (b_i - a_i)s_i$$

$$\implies s_j = \sum_{i=1, i \neq j}^k \frac{b_i - a_i}{a_j - b_j} s_i \quad \text{since } (a_j - b_j) \neq 0$$

And we have found an element of S that is a linear combination of other elements of S .

$$S' := S \setminus \{s_j\} \subset S, \text{span}(S') = V$$

This contradicts the minimality of S . Hence $a_i = b_i$ for all i .

(3) \implies (4) WTS unique representation implies maximal linear independence.

Since $0 \cdot s_1 + 0 \cdot s_2 + \dots + 0 \cdot s_k = 0$, and representations are unique,

$$a_1 s_1 + a_2 s_2 + \dots + a_k s_k \implies a_1 = a_2 = \dots = 0$$

Hence S is linearly independent.

To show S is maximally linearly independent, take any $v \in V \setminus S$. By hypothesis, (assuming (3))

$$v = a_1 s_1 + a_2 s_2 + \dots + a_k s_k$$

Hence,

$$a_1 s_1 + a_2 s_2 + \dots + a_k s_k - v = 0$$

Therefore, $S \cup \{v\}$ is not linearly independent.

(4) \implies (1). WTS that maximal linear independence implies S is a basis.

It suffices to show that $\text{span}(S) = V$. Assume towards a contradiction otherwise, then $\text{span}(S) \neq V, \exists v \in V \setminus \text{span}(S)$. By lemma,

$$\bar{S} = S \cup \{v\}$$

is also linearly independent. $S \subset \bar{S}$. This contradicts the assumption that S is maximally linearly independent. \checkmark

Corollary 5.4. Every finitely generated vector space V has a basis.

Proof. Since V is finitely generated, we can find $S \subseteq V$ finite s.t. $\text{span}(S) = V$.

We can successively remove elements from S until it is a minimal set of generators. \checkmark

Remark. Any vector space has a basis.

5.2 Dimension

Lemma 5.5. (Exchange Lemma): Let V be a F -vector space with basis $S = \{s_1, \dots, s_n\}$. Let w be

$$w = a_1 s_1 + \dots + a_n s_n$$

If k is such that $a_k \neq 0$, then

$$S' := \{s_1, \dots, s_{k-1}, w, s_{k+1}, \dots, s_n\}$$

is also a basis.

Proof. WLOG assume $a_1 \neq 0$. $S' = \{w, s_2, \dots, s_n\}$.

(1) WTS that $\text{span}(S') = \text{span}(S) = V$.

Since $a_1 \neq 0$,

$$\begin{aligned} w &= a_1 s_1 + \dots + a_n s_n \\ s_1 &= \frac{1}{a_1} w - \frac{a_2}{a_1} s_2 - \frac{a_3}{a_1} s_3 - \dots - \frac{a_n}{a_1} s_n \in \text{span}(S') \end{aligned}$$

Hence

$$S \subseteq \text{span}(S') \implies V \subseteq \text{span}(S')$$

also

$$\text{span}(S') \subseteq V \implies \text{span}(S') \subseteq V$$

Hence $V = \text{span}(S')$.

(2) WTS that S' linearly independent.

Take $c, c_2, \dots, c_n \in F$ so that

$$cw + c_2 s_2 + \dots + c_n s_n = 0$$

Since $w = a_1 s_1 + \dots + a_n s_n$, substituting, we get

$$ca_1 s_1 + (ca_2 + c_2) s_2 + \dots + (ca_n + c_n) s_n = 0$$

By linearly independence of S ,

$$ca_1 = (ca_2 + c_2) = \dots = (ca_n + c_n) = 0$$

Hence

$$c = c_2 = \dots = c_n = 0$$

\checkmark

Theorem 5.6. (Exchange Theorem): Let V be a F -vector space with basis $S = \{s_1, \dots, s_n\}$. Let $T = \{t_1, t_2, \dots, t_m\}$ be a linear independent subset of V . Then $m \leq n$ and there are m elements in S which can be exchanged with elements of T to obtain a new basis, i.e. we can form

$$\{t_1, t_2, \dots, t_m, s_{m+1}, \dots, s_n\}$$

Proof.

By induction in m .

Case $m = 0$ is immediate.

Assume that $m \geq 1$ and that the Exchange Theorem is true for $m - 1$. Let $T = \{t_1, \dots, t_m\}$. $T_0 = \{t_1, \dots, t_{m-1}\}$ is linearly independent as well.

By induction hypothesis, $m - 1 \leq n$ and after relabelling, S is $\{t_1, \dots, t_{m-1}, s_m, s_{m+1}, \dots, s_n\}$.

(1) We want to show that $m \leq n$. Since we assume that induction hypothesis is true, $m - 1 \leq n$. This implies either $m = n + 1$ or $m \leq n$.

If $m - 1 = n$, then $\{t_1, \dots, t_{m-1}\}$ is a new basis. However, $\{t_1, \dots, t_m\}$ is linearly independent. This contradicts with the fact that basis are maximally linearly independent. Hence $m = n$

(2) Since $\{t_1, \dots, t_{m-1}, s_m, \dots, s_n\}$ is a basis, we can write

$$t_m = \sum_{i=1}^{m-1} a_i t_i + \sum_{i=m}^n a_i s_i$$

Rearranging, we get

$$a_1 t_1 + \dots + a_{m-1} t_{m-1} - t_m = -a_m s_m - \dots - a_n s_n$$

Since $\{t_1, \dots, t_m\}$ is linearly independent, the LHS is non-zero, and there must be some $a_k, m \leq k \leq n$ such that $a_k \neq 0$.

By exchange lemma, in the basis $\{t_1, \dots, t_{m-1}, s_m, \dots, s_n\}$, we can replace s_k with t_m , to get a new basis

$$S \setminus \{s_k\} \cup \{t_m\}$$

✓

Corollary 5.7. (Basis extension theorem): Let V be a finitely-generated F -vector space. Every linearly independent set $\{t_1, \dots, t_m\}$ can be extended to form a basis for V . I.e. we can find

$$t_{m+1}, \dots, t_n \in V \text{ such that } S = \{t_1, \dots, t_m, t_{m+1}, \dots, t_n\}, n \geq m$$

Proof. By exchange theorem, consider any basis S . T is a linearly independent set. We can choose t_{m+1}, \dots, t_n to be s_{m+1}, \dots, s_n respectively. ✓

6 Class 6

6.1 Basis, cont'd

Corollary 6.1. (Bases have equan cardinality): If V has a finite basis of n elements, then any other basis of V is finite with exactly n elements.

Proof. Let $S = \{s_1, \dots, s_n\}$ be a basis of V with n elements.

Any other basis has to be finite. Otherwise, we would have an infinitely linearly independent set. In particular, we can find $n + 1$ linearly independent vectors, which contradicts the exchange theorem.

If anther basis has k elements, by exchange theorem, taking the other basis to be the linearly independent set, $k \leq n$. Also by exchange theorem, $n \leq k$. Hence $n = k$. ✓

Definition 6.2. (Dimension): Let V be a F -vector space over V . Then

$$\dim V = \begin{cases} \infty & \text{if } V \text{ not finitely generated} \\ n & \text{if } V \text{ has a basis of } n \text{ elements} \end{cases}$$

Remark. "finitely generated" means "finite dimensional". Henceforth we will use "finite dimensional".

Remark. $\dim F^n = n$, because $\{e_1, \dots, e_n\}$ is a basis.

Corollary 6.3. Let V be a finite-dimensional F -vector space $W < V$ is a proper subspace (i.e. $W \leq V, W \neq V$), then $\dim W < \dim V$

Proof. Let $n = \dim V$. We can't have more than n linearly independent vectors in V . Hence $\dim W < \infty$.

Let $m = \dim W$, and $\{w_1, \dots, w_n\}$ be a basis for W . Since $W \subset V$, there is $u \in V \setminus \{W\}$.

$$u \notin \text{span}(w_1, \dots, w_n)$$

Hence w_1, \dots, w_n, u is linearly indepdent.

$$\dim V \geq m + 1 > m = \dim W$$

✓

Theorem 6.4. (Dimension of sum of subspaces): Let V be a finite-dimensional F -vector space. Let W_1, W_2 be subspaces of V . Then

1. $\dim(W_1 + W_2) = \dim W_1 + \dim W_2 - \dim(W_1 \cap W_2)$
2. If $W_1 \cap W_2 = \{0\}$, then $\dim(W_1 \oplus W_2) = \dim W_1 + \dim W_2$

Proof.

(1) \implies (2): \emptyset is a basis of $\{0\}$, so $\dim\{0\} = 0$.

(1): Let $d_0 = \dim(W_1 \cap W_2)$, $d_1 = \dim W_1$, $d_2 = \dim W_2$. Let $T = \{t_1, t_2, \dots, t_{d_0}\}$ be a basis for $W_1 \cap W_2$. Complete T to be a basis of W_1 and W_2 .

$$\begin{aligned} \beta_{W_1} &= T \cup S, S = \{s_1, \dots, s_{d_1-d_0}\} \\ \beta_{W_2} &= T \cup R, R = \{r_1, \dots, r_{d_2-d_0}\} \end{aligned}$$

Claim: $\beta = T \cup S \cup R$ is a basis for $W_1 + W_2$.

If claim were true, then

$$\begin{aligned} \dim(W_1 + W_2) &= |T| + |S| + |R| \\ &= d_0 + (d_1 - d_0) + (d_2 - d_0) \\ &= d_1 + d_2 - d_0 \\ &= \dim W_1 + \dim W_2 - \dim(W_1 \cap W_2) \end{aligned}$$

WTS $\langle T \cup S \cup R \rangle$ spanning:

Since $\langle T \cup S \rangle = W_1$, $\langle T \cup R \rangle = W_2$,

$$W_1 + W_2 \subseteq \langle T \cup S \cup R \rangle$$

We also have $\langle T \cup S \cup R \rangle \subseteq W_1 + W_2$. Hence

$$\langle T \cup S \cup R \rangle = W_1 + W_2$$

WTS $(T \cup S \cup R)$ linearly independent:

Suppose

$$\begin{aligned} 0 &= \sum_{i=1}^{d_0} a_i t_i + \sum_{j=1}^{d_1-d_0} b_j s_j + \sum_{k=1}^{d_2-d_0} c_k r_k \\ &= v_0 + v_1 + v_2 \end{aligned}$$

Then

$$v_0 + v_1 = -v_2 \in W_1 \cap W_2$$

Since $v_0 \in W_1 \cap W_2, v_1 \in W_1, (v_0 + v_1) \in W_1, -v_2 \in W_2$.

Since $v_0 + v_1 \in W_1 \cap W_2$, we can express it in terms of the basis

$$v_0 + v_1 = -v_2 = \sum_{i=1}^{d_0} \lambda_i t_i = \sum_{i=1}^{d_0} a_i t_i + \sum_{j=1}^{d_1-d_0} b_j s_j$$

Since $T \cup S$ is a basis for W_1 , by the fact that representations are unique, we know that all $b_j = 0$.

Now we have

$$0 = v_0 + v_2 = \sum_{i=1}^{d_0} a_i t_i + \sum_{k=1}^{d_2-d_0} c_k r_k$$

Since $T \cup R$ is a basis for W_2 , $a_i = c_k = 0$ for all i, k .

✓

7 Class 7

7.1 Matrices and Systems of linear equations

Definition 7.1. (Matrix): A $m \times n$ matrix over field F is an array of elements $a_{ij} \in F$ of the form

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

Where m is the number of rows and n is the number of columns.

We denote $Mat_{m \times n}(F)$ the set of all such matrices, or $F^{m \times n}$.

. A_{ij} denotes the (i, j) entry of matrix $A \in Mat_{m \times n}(F)$.

Remark. $F^{m \times n}$ is a vector space with sum and scalar multiplication defined entrywise.

Remark. $\dim F^{m \times n} = mn$.

Proof. We present a basis with mn elements. Consider

$$\{E^{ij}\}_{1 \leq i \leq m, 1 \leq j \leq n}$$

Where

$$(E^{ij})_{kl} = \begin{cases} 1 & \text{if } (k, l) = (i, j) \\ 0 & \text{otherwise} \end{cases}$$

✓

Definition 7.2. (Matrix Multiplication): $A \in F^{m \times n}, B \in F^{n \times r}$. Then, $AB \in F^{m \times r}$ is defined by

$$(AB)_{ij} = \sum_{k=1}^n A_{ik} B_{kj}$$

I.e. the (i, j) -th entry of AB is the dot product of the i -th row of A with the j -th column of B .

Remark. Properties of matrix multiplication

- In general, for $A, B \in F^{n \times m}$, $AB \neq BA$
- $A \in F^{m \times n}, B \in F^{n \times r}, C \in F^{r \times s}$, $(AB)C = A(BC)$.

Definition 7.3. (Systems of linear equations): Let $b_1, b_2, \dots, b_n \in F, a_{ij} \in F, \forall 1 \leq i \leq m, 1 \leq j \leq n$, the set of equations

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m \end{cases}$$

is called a system of m -linear equations in n unknowns.

Remark. In matrix notation, let A, B

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix} \in F^{m \times n}$$

$$b = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix} \in F^{m \times 1}$$

The system of m -linear equations in n variables is denoted

$$Ax = b$$

Where

$$x = \begin{pmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{pmatrix} \in F^{n \times 1}$$

Definition 7.4. (Homogeneity): A system $Ax = b$ is homogenous if $b = 0 \in F^n$. Otherwise it is inhomogenous.

Remark. A homogenous system has at least one solution with $x = 0$. Otherwise, this is not guaranteed.

Definition 7.5. (Solution set): The solution set of a linear system $Ax = b$ is the set of elements in $F^{n \times 1}$ such that $Ax = b$

$$\{x \in F^{n \times 1} : Ax = b\}$$

Remark. If the system is homogenous, then the solution set is a subspace.

7.2 Echelon form and Row-reduced echelon form

Definition 7.6. (Echelon form): $A \in F^{m \times n}$ is in echelon form if

1. There exists some $r, 1 \leq r \leq m$ so that every row of index less than or equal to r has at least 1 non-zero entry, and every row of index greater than r is zero
2. for every $i \leq r$, consider the lowest index j_i that has a non-zero entry, i.e.

$$j_i := \min\{1 \leq j \leq n : a_{ji} \neq 0\}$$

Then

$$a_{ij_i} = 1$$

3. $j_1 \leq j_2 \leq j_3 \dots < j_r$

Remark. The a_{ij_i} are referred to as pivots.

- If A is in echelon form, then we can find the solution set.
- By relabelling the variables, assume we have pivots in the first r columns, $Ax = b$ becomes

$$\left(\begin{array}{cccc|c} 1 & & & & b_1 \\ 0 & 1 & & & b_2 \\ & & \ddots & & \vdots \\ 0 & & & 1 & b_r \\ \hline 0 & 0 & \dots & 0 & b_{r+1} \\ 0 & 0 & \dots & 0 & \vdots \\ 0 & 0 & \dots & 0 & b_m \end{array} \right)$$

- If there is some $i > r$ for which $b_i \neq 0$, then there is no solution.
- If all $b_i = 0$ for $i > r$, the variables x_1, x_2, \dots, x_r can be solved in terms of the variables $x_{r+1}, x_{r+2}, \dots, x_n$

Definition 7.7. (Row-reduced echelon form): A is in the row-reduced echelon form if A is in the echelon form and all entries above the pivots are zero.

Definition 7.8. (Elementary row operations):

- **RO1:** Exchange 2 different rows
- **RO2:** Add λ times i -th row to the j -th row where $\lambda \in F \setminus \{0\}, i \neq j$ and replacing row j with the result
- **RO3:** Multiply a row by a non-zero scalar in F

Theorem 7.9. (Row-reduced echelon form):

1. Every matrix A can be put into row-reduced echelon form using finitely many elementary row operations
2. If $Ax = b$ is a system of linear equations and $(\tilde{A}|\tilde{b})$ is the matrix obtained from $(A|b)$ by performing the row operations that **put A in row-reduced echelon form**, then they have the same solution set

Remark. $(A|b)$ denotes the $m \times (n + 1)$ matrix obtained from A by appending $b \in F^{m \times 1}$ to $A \in F^{m \times n}$.

Proof.

(1): Assume $A \in F^{m \times n}$, $A \neq 0$, find the first non-zero column of A ,

$$j_1 := \min\{1 \leq j \leq n : a_{ij} \neq 0 \text{ for some } i\}$$

- If $A_{1j_1} \neq 0$, multiply the first row by $(A_{1j_1})^{-1}$ (RO3), i.e. *creating a pivot in the first row in the $(1, j_1)$ position*. We can make every other entry of that column 0 (finite number of RO2).
- If $A_{1j_1} = 0$, let $i_1 \neq 1$ be the first non-zero entry in the j_1 column and exchange row 1 with row i_1 (RO1)

$$\begin{pmatrix} 0 & \cdots & 0 & 1 & \cdots & \cdots & \cdots \\ 0 & \cdots & 0 & 0 & & & \\ \vdots & & \vdots & \vdots & & & \\ 0 & \cdots & 0 & 0 & & A_2 & \end{pmatrix}$$

Repeat the process with A_2 to get the result after finitely many steps. Finally, we use RO2 to convert the matrix from echelon form to row-reduced echelon form.

(2): It suffices to show that each elementary row operation does not change the solution set. RO1 and RO3 are obvious.

For RO2, let

$$(1) \begin{cases} a_{i1}x_1 + a_{i2}x_2 + \cdots + a_{in}x_n = b_i \\ a_{j1}x_1 + a_{j2}x_2 + \cdots + a_{jn}x_n = b_j \end{cases}$$

$$(2) \begin{cases} a_{i1}x_1 + a_{i2}x_2 + \cdots + a_{in}x_n = b_i \\ (a_{j1} + a_{i1})x_1 + (a_{j2} + a_{i2})x_2 + \cdots + (a_{jn} + a_{in})x_n = b_j \end{cases}$$

Suppose \mathbf{x} satisfies (1), add $\lambda 1.1$ to 1.2, then 2.2 holds. Hence \mathbf{x} is also a solution for (2). Likewise, if \mathbf{x} is a solution to (2), do $2.2 - \lambda 1.1$, then 1.2 also holds.

✓

Corollary 7.10. If $A \in F^{m \times n}$ and $m < n$ then $Ax = 0$ has a non-trivial solution.

Proof. Let \tilde{A} be the row-reduced echelon form of A , then by theorem above,

$$Ax = 0 \Leftrightarrow \tilde{A}x = 0$$

The matrix \tilde{A} has $0 \leq r \leq m$ non-zero rows which corresponds to the number of pivots, which is the number of non-free variables. \tilde{A} has $n - r$ free variables

$$\begin{aligned} r &\leq m \\ -r &\geq -m \\ n - r &\geq n - m > 0 \end{aligned}$$

$\tilde{A}x = 0$ has a non-trivial solution by taking all free variables say 1.

✓

Corollary 7.11. Let $A \in F^{n \times n}$ and \tilde{A} be the row-reduced echelon form of A . Then, \tilde{A} is the identity if and only if $x = 0$ is the unique solution to $Ax = 0$.

Proof.

(\Rightarrow):

$$\begin{aligned} \tilde{A} = I &\Rightarrow Ax = 0 \Leftrightarrow \tilde{A}x = 0 \\ &\Leftrightarrow Ix = 0 \\ &\Leftrightarrow x = 0 \end{aligned}$$

(\Leftarrow): Assume $x = 0$ is the only solution to $Ax = 0$. Then \tilde{A} does not have free variables, $r \geq n$. However, $r \leq n$ always. Hence $r = n$. Therefore $\tilde{A} = I$.

✓

8 Class 8

8.1 Elementary Matrices and Invertible Matrices

Definition 8.1. (Elementary matrix) An elementary matrix is a matrix that can be obtained from the identity matrix by a single elementary row operation.

Example. In \mathbb{R}^2 , the following are elementary matrices

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ a & 1 \end{pmatrix}, \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix},$$

for $a \in \mathbb{R}, a \neq 0$

Theorem 8.2. Let e be an elementary row operation and let $E = e(I)$ be the corresponding matrix of size $m \times m$.

Then $e(A) = EA$ for every $m \times n$ matrix A

Proof. RO1:

RO2: replace row r by row $r + c \times$ row r .

$$E_{ik} = \begin{cases} \delta_{ik}, i \neq r \\ \delta_{rk} + c + \delta_{sk}, i = r \end{cases}$$

Then

$$(EA)_{ij} = \sum_{k=1}^m E_{ik} A_{kj} = \begin{cases} A_{ik}, i \neq r, A_{rj} + cA_{sj}, i = r \end{cases}$$

RO3:

✓

Example. Let e be the row operation of adding 2 times the first row to the second row, and

$$A = \begin{pmatrix} 2 & 3 & 4 & 5 \\ 1 & 1 & 0 & 1 \end{pmatrix}$$

$$e(A) = \begin{pmatrix} 2 & 3 & 4 & 5 \\ 5 & 7 & 8 & 11 \end{pmatrix}$$

Also,

$$E = e(I) = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$$

$$EA = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 2 & 3 & 4 & 5 \\ 1 & 1 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 3 & 4 & 5 \\ 5 & 7 & 8 & 11 \end{pmatrix}$$

Corollary 8.3. Let $A, B \in F^{m \times n}$, A can be transformed into B by a finite series of elementary matrices if and only if $B = PA$, where P is some product of elementary matrices.

Proof. \implies : If one can take A into B with row operations e_1, e_2, \dots, e_k , in this order, let $E_i = e_i(I)$, then

$$B = E_k E_{k-1} E_{k-2} \dots E_1 A$$

Take

$$P = E_k E_{k-1} E_{k-2} \dots E_1$$

\Leftarrow Let $B = E_k E_{k-1} \dots E_1 A$. Define

$$e_i(A) := E_i A$$

We can follow the row operations dictated by the E_i 's to get from A to B .

✓

Definition 8.4. If A can be transformed into B by a series of finitely many row operations, then so can B be transformed into A (i.e. row operations can be reversed), and A and B are called row equivalent matrices.

Definition 8.5. Let $A \in \text{Matr}_n(F)$

9 Class 9

10 Class 10

11 Class 11

12 Class 12

13 Class 13

14 Class 14

14.1 Quotients, cont'd

Recall from last time (homomorphism theorem) that if $\varphi : V \rightarrow W$ is a linear map between F -vector spaces, then

$$\tilde{\varphi} : V/\ker \varphi \rightarrow \operatorname{Im} \varphi, [v] \mapsto \varphi(v)$$

is well defined isomorphism.

Corollary 14.1. Every linear map $\varphi : V \rightarrow W$ factors as

$$\varphi = i \circ \bar{\varphi} \circ \pi$$

where

- $\pi : V \rightarrow V/\ker \varphi$ is the canonical projection
- $i : \operatorname{Im} \varphi \rightarrow W$ is the inclusion map
- $\bar{\varphi} : V/\ker \varphi \rightarrow \operatorname{Im} \varphi$ is isomorphism

Proposition 14.2. (Dimension of a quotient space) Let V be a finite dimensional vector space over F , and let $W \leq V$, then

$$\dim(V/W) = \dim V - \dim W$$

Proof. Say $\dim W = m$. Take (w_1, \dots, w_m) basis for W . Extend it to a basis of V , $S = (w_1, w_2, \dots, w_m, v_{m+1}, v_{m+2}, \dots, v_n)$ basis of V .

WTS that $([v_{m+1}], [v_{m+2}], \dots, [v_n])$ is a basis for V/W .

Let $v \in V$ Since S is a basis for V

$$\begin{aligned} v &= a_1 w_1 + \dots + a_m w_m + a_{m+1} v_{m+1} + \dots + a_n v_n \\ \implies [v] &= [a_1 w_1 + \dots + a_m w_m] + [a_{m+1} v_{m+1} + \dots + a_n v_n] \end{aligned}$$

$$\implies [v] = [0] + a_{m+1} [v_{m+1}] + \dots + a_n [v_n]$$

Hence $[v_{m+1}], \dots, [v_n]$ spans V/W .

To show linear independence, let

$$b_{m+1} [v_{m+1}] + \dots + b_n [v_n] = 0$$

for some b_{m+1}, \dots, b_n .

$$\begin{aligned} b_{m+1} [v_{m+1}] + \dots + b_n [v_n] &= 0 \\ \implies \left[\sum_{i=m+1}^n b_i v_i \right] &= [0] \end{aligned}$$

That is,

$$\sum_{i=m+1}^n b_i v_i \in W$$

By linear independence of v_i 's in S ,

$$b_{m+1} = \dots = b_n = 0$$

Hence,

$$\begin{aligned} \dim(V/W) &= \#\{[v_{m+1}], \dots, [v_n]\} \\ &= n - m \\ &= \dim V - \dim W \end{aligned}$$

✓

Corollary 14.3. (New proof of dimension formula for linear maps)

Let $\varphi : V \rightarrow W$ be a linear map between F -vector spaces.

$$\dim V = \dim \ker \varphi + \dim \operatorname{Im} \varphi$$

Proof. By the homomorphism theorem,

$$\dim V/\ker \varphi \cong \operatorname{Im} \varphi$$

Then

$$\dim V / \ker \varphi = \dim \operatorname{Im} \varphi \text{ by Homomorphism Theorem}$$

$$\dim V \ker \varphi = \dim V - \dim \ker \varphi \text{ by above proposition}$$

Hence

$$\dim V = \dim \ker \varphi + \dim \operatorname{Im} \varphi$$

✓

Example. (Quotient capturing Taylor expansion)

Let $V = C^\infty[-1, 1]$ be the space of smooth real-valued functions on $[-1, 1]$ and fix $d \in \mathbb{N}_{\geq 0}$.

Define

$$W_d = \{f \in C^\infty[-1, 1] \text{ s.t. } f^{(k)}(0) = 0, k = 0, 1, 2, \dots, d\} \leq V$$

W_d consists of functions whose Taylor polynomial of degree d at 0 vanishes completely.

Then the quotient

$$V/W_d$$

is naturally isomorphic to the space of polynomials of degree at most d .

The isomorphism is induced by the map

$$\Phi : C^\infty[-1, 1] \rightarrow \mathcal{P}_d, f \mapsto \Phi(f)(x) = f(0) + f'(0)x + \frac{1}{2!}f''(0)x^2 + \dots + \frac{1}{d!}f^{(d)}(0)x^d$$

One has

$$V/W_d = V/\ker \Phi \cong \operatorname{Im} \Phi = \mathcal{P}_d$$

Example. Recall $V = \mathbb{R}^2, W = \operatorname{span}(1, 0)$.

Now, we know that

$$\dim V/W = \dim \mathbb{R}^2 - \dim W = 1$$

14.2 Linear Functionals

14.2.1 Dual space

Definition 14.4. (Linear Functionals) Let V be an F -VS. A linear map $f : V \rightarrow F$ is also called a **linear functional**.

Definition 14.5. Let F be a field and V be a F -vector space. The dual space is defined as

$$V^* := \operatorname{Hom}_F(V, F)$$

i.e. the vector space of all linear functionals on V .

Example. Examples of linear functionals

- sum of constants of polynomial Let $V = \mathcal{P}_d(\mathbb{R})$, then

$$f : \mathcal{P}_d(\mathbb{R}) \rightarrow \mathbb{R}, a_0 + a_1x + \dots + a_dx^d \mapsto a_0 + a_1 + \dots + a_d$$

- evaluation map Let $V = C^0[-1, 1]$, then

$$F_0 : C^0[-1, 1] \rightarrow \mathbb{R}, g \mapsto g(0)$$

- integration map

$$\Phi : C[a, b] \rightarrow \mathbb{R}, f \mapsto \int_a^b f(x)dx$$

- linear functional in F^n Fix $a_1, a_2, \dots, a_n \in F$, define

$$f : F^n \rightarrow F, \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} \mapsto a_1v_1 + \dots + a_nv_n$$

Counter examples of linear functionals

- finding the length

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}, (x, y, z) \mapsto \sqrt{x^2 + y^2 + z^2}$$

is not a linear functional.

$$f(-(1, 0, 0)) \neq -f(1, 0, 0)$$

- product of coordinates

$$F : \mathbb{R}^2 \rightarrow \mathbb{R}, (x, y) \mapsto xy$$

is not a linear functional. Take $v_1 = (1, 0), v_2 = (0, 1)$

$$F(v_1) = F(v_2) = 0$$

$$F(v_1) + F(v_2) = 0 \neq F(v_1 + v_2) = 1$$

Remark. Every linear functional in F^n has the form

$$f : F^n \rightarrow F, \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} \mapsto a_1 v_1 + \dots a_n v_n$$

Proof. Let $g \in (F^n)^*$, then

$$\begin{aligned} g(v) &= g \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} = g(v_1 e_1 + \dots v_n e_n) \\ &= v_1 g(e_1) + \dots v_n g(e_n) \text{ by linearity of } g. \end{aligned}$$

if you define $a_i = g(e_i), 1 \leq i \leq n$, then

$$g(v) = \sum_{i=1}^n a_i \pi_i$$

✓

Theorem 14.6. Let V be a vector space over F with basis $S = (s_1, s_2, \dots, s_n)$. Then

1. $\dim V^* = \dim V$
2. Let f_i be linear map such that

$$f_i(s_j) = \delta_{ij} = \begin{cases} 1 & \text{if } j = i \\ 0 & \text{otherwise} \end{cases}$$

Then $S^* = (f_1, f_2, \dots, f_n)$ is a basis for V^* .

Remark. Recall $\dim W = m, \dim V = n$,

$$\dim \text{Hom}_F(V, W) = mn$$

Proof.

Proof of (1):

$$\begin{aligned} \dim V^* &= \dim \text{Hom}_F(V, F) \\ &= \dim V \times \dim F \\ &= \dim V \end{aligned}$$

Proof of (2): since we know that $\dim V^* = n$, it suffices to show that $S^* = (f_1, f_2, \dots, f_n)$ linearly independent in V^* .

We take a linear combination of S^* that gives the 0 functional.

$$a_1 f_1 + a_2 f_2 + \dots + a_n f_n = 0$$

Apply functionals at s_j

$$\begin{aligned} (a_1 f_1 + a_2 f_2 + \dots + a_n f_n)(s_j) &= 0(s_j) = 0 \\ \implies a_1 f_1(s_j) + a_2 f_2(s_j) + \dots + a_n f_n(s_j) &= 0 \\ \implies a_j f_j(s_j) &= 0 \\ \implies a_j &= 0 \end{aligned}$$

This is true for all $1 \leq j \leq n$, therefore $S^* = (f_1, f_2, \dots, f_n)$ linearly independent.

✓

Definition 14.7. $S^* = (f_1, f_2, \dots, f_n)$ from theorem above is called the dual basis of S .

Each f_i is denoted

$$f_i = S_i^*$$

Example. Let $V = F^n$, and $S = (e_1, e_2, \dots, e_n)$ is the standard basis where $e_i = (0, 0, \dots, 1, \dots, 0)^T$ (only nonzero element is 1 at the i -th position).

Then

$$\begin{aligned}
 e_i^* \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} &= e_i^* \left(\sum_{j=1}^n v_j e_j \right) \\
 &= \sum_{j=1}^n v_j e_i^*(e_j) \\
 &= v_i e_i^*(e_i) \\
 &= v_i
 \end{aligned}$$

14.2.2 Duality Theorem

Definition 14.8. Since V^* is again a vector space over F . Define the bidual space as

$$V^{**} := (V^*)^* = \text{Hom}_F(V^*, F)$$

Remark. If $\dim V < \infty$,

$$\dim(V^{**}) = \dim V^* = \dim V$$

Theorem 14.9. Let V be a finite-dimensional F -vector space. Then, there exists a natural isomorphism

$$\Theta : V \rightarrow V^{**} = \text{Hom}_F(V^*, F), v \mapsto \theta(v) = \theta_v$$

Where

$$\theta_v(f) = f(v) \text{ for all } f \in V^*$$

i.e. θ_v is an evaluation functional (taking linear functionals to scalars).