

Important Notice to Authors

No further publication processing will occur until we receive your response to this proof.

Attached is a PDF proof of your forthcoming article in PRC. Your article has 15 pages and the Accession Code is **CM10581**.

Please note that as part of the production process, APS converts all articles, regardless of their original source, into standardized XML that in turn is used to create the PDF and online versions of the article as well as to populate third-party systems such as Portico, Crossref, and Web of Science. We share our authors' high expectations for the fidelity of the conversion into XML and for the accuracy and appearance of the final, formatted PDF. This process works exceptionally well for the vast majority of articles; however, please check carefully all key elements of your PDF proof, particularly any equations or tables.

Figures submitted electronically as separate files containing color appear in color in the online journal. However, all figures will appear as grayscale images in the print journal unless the color figure charges have been paid in advance, in accordance with our policy for color in print (<https://journals.aps.org/authors/color-figures-print>).

Specific Questions and Comments to Address for This Paper

FQ: This funding provider could not be uniquely identified during our search of the FundRef registry (or no Contract or Grant number was detected). Please check information and amend if incomplete or incorrect.

Open Funder Registry: Information about an article's funding sources is now submitted to Crossref to help you comply with current or future funding agency mandates. Crossref's Open Funder Registry (<https://www.crossref.org/services/funder-registry/>) is the definitive registry of funding agencies. Please ensure that your acknowledgments include all sources of funding for your article following any requirements of your funding sources. Where possible, please include grant and award ids. Please carefully check the following funder information we have already extracted from your article and ensure its accuracy and completeness:

Other Items to Check

- Please note that the original manuscript has been converted to XML prior to the creation of the PDF proof, as described above. Please carefully check all key elements of the paper, particularly the equations and tabular data.
- Title: Please check; be mindful that the title may have been changed during the peer-review process.
- Author list: Please make sure all authors are presented, in the appropriate order, and that all names are spelled correctly.
- Please make sure you have inserted a byline footnote containing the email address for the corresponding author, if desired. Please note that this is not inserted automatically by this journal.
- Affiliations: Please check to be sure the institution names are spelled correctly and attributed to the appropriate author(s).
- Receipt date: Please confirm accuracy.
- Acknowledgments: Please be sure to appropriately acknowledge all funding sources.
- Hyphenation: Please note hyphens may have been inserted in word pairs that function as adjectives when they occur before a noun, as in "x-ray diffraction," "4-mm-long gas cell," and "R-matrix theory." However, hyphens are deleted from word pairs when they are not used as adjectives before nouns, as in "emission by x rays," "was 4 mm in length," and "the R matrix is tested."

Note also that Physical Review follows U.S. English guidelines in that hyphens are not used after prefixes or before suffixes: superresolution, quasiequilibrium, nanoprecipitates, resonancelike, clockwise.

- Please check that your figures are accurate and sized properly. Make sure all labeling is sufficiently legible. Figure quality in this proof is representative of the quality to be used in the online journal. To achieve manageable file size for online delivery, some compression and downsampling of figures may have occurred. Fine details may have become somewhat fuzzy, especially in color figures. The print journal uses files of higher resolution and therefore details may be sharper in print. Figures to be published in color online will appear in color on these proofs if viewed on a color monitor or printed on a color printer.
- Please check to ensure that reference titles are given as appropriate.
- Overall, please proofread the entire *formatted* article very carefully. The redlined PDF should be used as a guide to see changes that were made during copyediting. However, note that some changes to math and/or layout may not be indicated.

Ways to Respond

- **Web:** If you accessed this proof online, follow the instructions on the web page to submit corrections.
- **Email:** Send corrections to prcproofs@aptaracorp.com
Subject: **CM10581** proof corrections
- **Fax:** Return this proof with corrections to +1.703.791.1217. Write **Attention:** PRC Project Manager and the Article ID, **CM10581**, on the proof copy unless it is already printed on your proof printout.

Comparative study of the requantization of the time-dependent mean field for the dynamics of nuclear pairing

Fang Ni¹ and Takashi Nakatsukasa^{2,1,3}

¹*Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba 305-8571, Japan*

²*Center for Computational Sciences, University of Tsukuba, Tsukuba 305-8577, Japan*

³*THES Research Group, RIKEN, Wako 351-0198, Japan*



(Received 11 December 2017; published xxxxxx)

To describe quantal collective phenomena, it is useful to requantize the time-dependent mean-field dynamics. We study the time-dependent Hartree-Fock-Bogoliubov (TDHFB) theory for the two-level pairing Hamiltonian, and compare results of different quantization methods. The one constructing microscopic wave functions, using the TDHFB trajectories fulfilling the Einstein-Brillouin-Keller quantization condition, turns out to be the most accurate. The method is based on the stationary-phase approximation to the path integral. We also examine the performance of the collective model which assumes that the pairing gap parameter is the collective coordinate. The applicability of the collective model is limited for the nuclear pairing with a small number of single-particle levels, because the pairing gap parameter represents only a half of the pairing collective space.

DOI: [10.1103/PhysRevC.00.004300](https://doi.org/10.1103/PhysRevC.00.004300)

I. INTRODUCTION

Pairing correlation plays a decisive role in a number of nuclear phenomena, which is especially important in open-shell nuclei. Many evidences of the pairing correlation were observed in experiment, including odd-even mass difference, moments of inertia of rotational bands, and quasiparticle spectra in odd nuclei. Even in closed-shell nuclei, the pairing dynamically plays an important role in elementary modes of excitation, such as pairing vibrations [1–3]. Properties of low-lying modes of excitation in even-even nuclei are expected to be determined dominantly by interplay between the pairing and the quadrupole correlations. However, the true nature of the low-lying excitations is still unclear, especially for excited $J^\pi = 0^+$ states [4,5]. Understanding the pairing dynamics is a key ingredient for solving mysteries of excited 0^+ states.

The ground states in many of the even-even nuclei are well described by the Bardeen-Cooper-Schrieffer (BCS) and the Hartree-Fock-Bogoliubov (HFB) theories [2,6]. Its time-dependent version, the time-dependent HFB (TDHFB) theory [7,8], is a natural extension of the static HFB theory. Thanks to increasing computational power, realistic applications of the TDHFB calculations in real time become available for studies of linear response properties [9–13] and of various nuclear dynamics [14–19]. The small-amplitude limit of the TDHFB is known to be the quasiparticle random phase approximation (QRPA). The QRPA was extensively utilized and successfully describes properties of giant resonances. Recently, the QRPA calculations with modern energy density functionals for giant resonances in deformed nuclei have become available [20–27]. In contrast, many of the low-lying excited states cannot be well reproduced by QRPA [8]. This may be because of their slowly moving large-amplitude nature.

In principle, the TDHFB dynamics can be applicable to large amplitude motion. The problem is that it is not easy

to determine quantum mechanical quantities, such as energy eigenvalues and transition matrix elements, from the TDHFB trajectories. In addition, the TDHFB lacks a part of quantum fluctuation associated with low-energy large amplitude collective motion, which leads to difficulties in the description of the quantum tunneling processes, such as spontaneous fission and sub-barrier fusion reaction. For this purpose, because the TDHF(B) trajectory corresponds to a stationary-phase limit of the path integral formulation, the requantization of the mean-field dynamics was proposed [28–33]. It recovers quantum fluctuations missing in the mean-field level, and possibly enables us to describe large-amplitude collective tunneling phenomena. The requantization of the TDHFB is particularly feasible for integrable systems, because the system is described by separable action-angle variables (I_k, ϕ_k) , leading to the Einstein-Brillouin-Keller (EBK) quantization condition. However, for nonintegrable systems in general, it is difficult to find suitable periodic orbits to quantize. A possible solution to this difficulty is to find a decoupled collective subspace spanned by a single coordinate and its conjugate momentum [8]. Because the one-dimensional system is integrable, the quantization is practicable.

Another somewhat phenomenological approach to nuclear collective dynamics is the collective model. In this approach, the collective Hamiltonian is constructed by choosing collective coordinates intuitively. The nuclear energy density functional model is often used as a tool for a microscopic derivation of the collective Hamiltonian [8]. The most well-known and successful model is the Bohr model [1], which was introduced to describe low-energy nuclear collective motion in quadrupole degrees of freedom with the deformation parameters (β, γ) and the Euler angles (ϕ, θ, ψ) . For the pairing motion, the collective coordinates are analogously chosen as the pair deformation (gap) Δ and the gauge angle Φ [34]. Based on the pairing collective Hamiltonian, effects of the pair motion

on the quadrupole vibrations have been discussed in literature [35,36]. However, it is not trivial whether the pair deformation is really a suitable choice for the collective coordinate, which should be investigated in a microscopic approach based on the TDHFB dynamics.

Our final goal is to study the role of large-amplitude pairing dynamics and to reveal the nature of the mysterious excited 0^+ states. As the first step, toward this goal, we investigate accuracy and applicability of the requantized TDHFB model for a two-level pairing model with equal degeneracy Ω [37], especially on the calculation of two-particle-transfer matrix elements. In Ref. [38], the two-particle-transfer matrix elements were evaluated as Fourier components of the time-dependent mean values of the pair-creation operators, which demonstrates nice agreement with the exact results at large Ω values ($\Omega = 40$). However, in realistic values of Ω , we will show that the deviation is substantial. The collective model of the pairing motion, which assumes the pair deformation as the collective coordinate, has a similar tendency, namely, applicability limited to large Ω cases [34]. This deficiency is mainly from the small collectivity in the pairing motion in realistic situations. In this paper, to improve the quantitative estimate of the matrix elements, we construct microscopic wave functions based on the EBK quantization for the integrable systems. The wave functions are obtained from the stationary-phase approximation for the path-integral form [39]. Its superiority to the other methods becomes more evident for smaller values of Ω .

The paper is organized as follows. In Sec. II, we derive a TDHFB classical Hamiltonian for the pairing model. In Sec. III, the requantization of the TDHFB is performed using different methods, based on the canonical and the EBK quantization. In Sec. IV, the numerical results for the two-level model are shown and compared with exact results. Properties of the pairing collective coordinate is also discussed. We give a conclusion in Sec. V.

II. CLASSICAL FORM OF TDHFB HAMILTONIAN

The Hamiltonian of the pairing model is given in terms of single-particle energies ϵ_l and the pairing strength g as

$$H = \sum_l \epsilon_l n_l - g \sum_{l,l'} S_l^+ S_{l'}^- \\ = \sum_l \epsilon_l (2S_l^0 + \Omega_l) - g S^+ S^-, \quad (2.1)$$

where we use the SU(2) quasispin operators, $S = \sum_l S_l$, with

$$S_l^0 = \frac{1}{2} \left(\sum_m a_{lm}^\dagger a_{lm} - \Omega_l \right), \quad (2.2)$$

$$S_l^+ = \sum_{m>0} a_{lm}^\dagger a_{l\bar{m}}^\dagger, \quad S_l^- = S_l^{+\dagger}. \quad (2.3)$$

Each single-particle energy ϵ_l possesses $(2\Omega_l)$ -fold degeneracy ($\Omega_l = j_l + 1/2$) and $\sum_{m>0}$ indicates the summation over $m = 1/2, 3/2, \dots$, and $\Omega_l - 1/2$. The occupation number of each level l is given by $n_l = \sum_m a_{lm}^\dagger a_{lm} = 2S_l^0 + \Omega_l$. The

quasispin operators satisfy the commutation relations,

$$[S_l^0, S_{l'}^\pm] = \pm \delta_{ll'} S_l^\pm, \quad [S_l^+, S_{l'}^-] = 2\delta_{ll'} S_l^0. \quad (2.4)$$

The magnitude of quasispin for each level is $S_l = \frac{1}{2}(\Omega_l - \nu_l)$, where ν_l is the seniority quantum number, namely the number of unpaired particles at each level l . In the present study, we only consider seniority zero states with $\nu = \sum_l \nu_l = 0$. The residual two-body interaction only consists of monopole pairing interaction which couples two particles to zero angular momentum. We obtain exact solutions either by solving the Richardson equation [40–42] or by diagonalizing the Hamiltonian using the quasispin symmetry.

A. Coherent-state representation of the TDHFB Hamiltonian

The coherent state for the seniority $\nu = 0$ states ($S_l = \Omega_l/2$) is constructed as

$$|Z(t)\rangle = \prod_l (1 + |Z_l(t)|^2)^{-\Omega_l/2} \exp[Z_l(t) S_l^+] |0\rangle, \quad (2.5)$$

where $|0\rangle$ is the vacuum (zero particle) state, $Z_l(t)$ are time-dependent complex variables which describe motion of the system. In the SU(2) quasispin representation, $|0\rangle = \prod_l |S_l, -S_l\rangle$. The coherent state $|Z(t)\rangle$ is a superposition of states with different particle numbers without unpaired particles. In the present pairing model, the coherent state is the same as the time-dependent BCS wave function with $Z_l(t) = v_l(t)/u_l(t)$, where $(u_l(t), v_l(t))$ are the time-dependent BCS u, v factors.

The TDHFB equation can be derived from the time-dependent variational principle,

$$\delta S = 0, \quad S \equiv \int \langle \phi(t) | i \frac{\partial}{\partial t} - H | \phi(t) \rangle dt, \quad (2.6)$$

where $|\phi(t)\rangle$ is the time-dependent generalized Slater determinant. In the present case, we adopt the coherent state of Eq. (2.5), $|\phi(t)\rangle = |Z(t)\rangle$. The action S is

$$S = \int \mathcal{L}(t) dt \\ = \int dt \left\{ \frac{i}{2} \sum_l \frac{\Omega_l}{1 + |Z_l|^2} (Z_l^* \dot{Z}_l - Z_l \dot{Z}_l^*) - \langle Z | H | Z \rangle \right\}, \quad (2.7)$$

and

$$\langle Z | H | Z \rangle = \sum_l \epsilon_l \frac{2\Omega_l |Z_l|^2}{1 + |Z_l|^2} - g \sum_l \frac{\Omega_l |Z_l|^2 (\Omega_l + |Z_l|^2)}{(1 + |Z_l|^2)^2} \\ - g \sum_{l_1 \neq l_2} \frac{\Omega_{l_1} Z_{l_1}}{1 + |Z_{l_1}|^2} \frac{\Omega_{l_2} Z_{l_2}^*}{1 + |Z_{l_2}|^2}. \quad (2.8)$$

We transform the complex variables Z_l into real variables (q_l, χ_l) by $Z_l = \tan \frac{\theta_l}{2} e^{-i\chi_l}$ and $q_l = \cos \theta_l$ ($0 \leq \theta \leq \pi$). The Lagrangian \mathcal{L} and the expectation value of Hamiltonian be-

163 come

$$\mathcal{L}(t) = \sum_l \frac{\Omega_l}{2} (1 - q_l) \dot{\chi}_l - \mathcal{H}(Z, Z^*), \quad (2.9)$$

$$\begin{aligned} \mathcal{H}(Z, Z^*) &\equiv \langle Z | H | Z \rangle \\ &= \sum_l \epsilon_l \Omega_l (1 - q_l) - \frac{g}{4} \\ &\quad \times \sum_l \Omega_l [\Omega_l (1 - q_l^2) + (1 - q_l)^2] \\ &\quad - \frac{g}{4} \sum_{l_1 \neq l_2} \Omega_{l_1} \Omega_{l_2} \sqrt{(1 - q_{l_1}^2)(1 - q_{l_2}^2)} e^{-i(\chi_{l_1} - \chi_{l_2})}. \end{aligned} \quad (2.10)$$

164 Here, χ_l represents a kind of gauge angle of each level, and
165 q_l are related to the occupation probability, $q_l = |u_l|^2 - |v_l|^2$.
166 If we choose χ_l as canonical coordinates, their conjugate
167 momenta are given by $p_l \equiv \partial \mathcal{H} / \partial \dot{\chi}_l = \Omega_l (1 - q_l) / 2$. Because
168 the Hamiltonian (2.10) depends only on the relative difference
169 in the gauge angles, the “global” gauge angle, $\Phi \propto \sum_l \chi_l$, is
170 a cyclic variable.

171 B. Two-level case

172 In a two-level system, it is convenient to define global and
173 relative gauge angles, Φ and ϕ , respectively.

$$\Phi \equiv \frac{\chi_1 + \chi_2}{2}, \quad \phi \equiv \chi_2 - \chi_1, \quad (2.11)$$

174 whose ranges are $0 \leq \Phi \leq 2\pi$ and $-2\pi \leq \phi \leq 2\pi$. Their
175 conjugate momenta (J, j) are given by

$$\begin{aligned} J &= \frac{\partial \mathcal{L}}{\partial \dot{\Phi}} = \sum_{l=1}^2 \frac{\Omega_l}{2} (1 - q_l), \\ j &= \frac{\partial \mathcal{L}}{\partial \dot{\phi}} = \frac{\Omega_2 (1 - q_2) - \Omega_1 (1 - q_1)}{4}. \end{aligned} \quad (2.12)$$

176 By calculating the occupation number n_l in the level l ,
177 the physical meaning of these conjugate momenta becomes
178 obvious:

$$n_l = \langle Z | n_l | Z \rangle = \Omega_l (1 - q_l). \quad (2.13)$$

179 Therefore, J corresponds to the total particle number $N =$
180 $\sum_l n_l$, while j corresponds to the difference of the occupation
181 number between the upper level and the lower level:

$$J = \frac{N}{2}, \quad j = \frac{n_2 - n_1}{4}. \quad (2.14)$$

182 The Hamiltonian in terms of these canonical variables
183 $(\phi, j; \Phi, J)$ is given by

$$\begin{aligned} \mathcal{H}(\phi, j; J) &= \sum_{l=1,2} \Omega_l \epsilon_l (1 - q_l) \\ &\quad - \frac{g}{4} \sum_{l=1,2} \Omega_l [\Omega_l (1 - q_l^2) + (1 - q_l)^2] \\ &\quad - \frac{g}{2} \Omega_1 \Omega_2 \sqrt{(1 - q_1^2)(1 - q_2^2)} \cos \phi, \end{aligned} \quad (2.15)$$

with

$$q_l = \frac{\Omega_l - J - 2(-1)^l j}{\Omega_l} \quad \text{for } l = 1, 2. \quad (2.16)$$

Note that the Hamiltonian does not depend on the global
gauge angle Φ . This leads to the particle number conservation,
 $dN/dt = 0$.

The TDHFB equation can be written in a form of the
classical equations of motion:

$$\frac{d\Phi}{dt} = \frac{\partial \mathcal{H}}{\partial J}, \quad \frac{dJ}{dt} = -\frac{\partial \mathcal{H}}{\partial \Phi}, \quad (2.17)$$

$$\frac{d\phi}{dt} = \frac{\partial \mathcal{H}}{\partial j}, \quad \frac{dj}{dt} = -\frac{\partial \mathcal{H}}{\partial \phi}. \quad (2.18)$$

Because the $J = N/2$ and the total energy E are constants
of motion, the TDHFB trajectories with given N and E are
determined in the two-dimensional phase space (ϕ, j) with the
condition,

$$\mathcal{H}(\phi(t), j(t); J = N/2) = E. \quad (2.19)$$

Examples of the classical trajectories in the phase space (ϕ, j)
are shown in Fig. 1. The figure shows contour lines of energy
for systems, which correspond to the TDHFB trajectories, with
 $\Omega_1 = \Omega_2 = 8$ and $N = 16$ with different values of g . The
transition from the normal to the superfluid phase takes place
at $g = \Delta \epsilon (2\Omega)^{-1}$ with $\Delta \epsilon = \epsilon_2 - \epsilon_1$. Using a dimensionless
parameter $x = 2g\Omega/\Delta \epsilon$, at $x < 1$, the ground state is normal
with the fully occupied lower level $n_1 = N$ and the empty
upper level ($n_2 = 0$). All the TDHFB trajectories represent
the rotational behavior with respect to the relative angle ϕ .
Here, the “rotational behavior” means that the motion spans
the whole region of the angle ϕ , while we use “vibrational” for
the classical motion in a bound region of ϕ ($-\pi < \phi < \pi$).
At $g = x = 0$, the Hamiltonian becomes independent from
 (Φ, ϕ) , then, the occupation numbers, n_1 and n_2 , are constants
of motion. At $x > 1$, the energy-minimum point and the closed
trajectories appear around $j = j_0$ ($-N/4 < j_0 < N/4$) and
 $\phi = 0$, which suggests the vibrational behavior for ϕ . At higher
energies, the trajectories become open (rotational-like), which
suggests a phase transition from super to normal phases as a
function of excitation energy.

In the single- j model, because the second term in the Hamil-
tonian (2.15) is absent, the Hamiltonian is exactly quadratic
with respect to N , $\mathcal{H}(N) = \mathcal{H}_0 + (N - N_0)^2 / (2\mathcal{J})$. The mo-
ment of inertia for the pair rotation is $\mathcal{J} = 2/g \times \Omega / (\Omega -$
 $1) \approx 2/g$ at $\Omega \rightarrow \infty$. In the multi- j case, the Hamiltonian
contains higher-order terms in general.

221 III. REQUANTIZATION OF TDHFB FOR THE 222 TWO-LEVEL PAIR MODEL

To determine energy eigenstates, we need to requantize the
TDHFB trajectories. Because the present two-level model is
integrable (Sec. II), it is feasible to apply the stationary-phase
approximation to the path integral expression. In addition, we
also study the canonical quantization of the TDHFB Hamilto-
nian, and the matrix elements extracted by the Fourier com-
ponents [38]. We show results of these different approaches to
the pairing model.




FIG. 1. Energy contour plot for $\Omega_1 = \Omega_2 = 8$ and $N = 16$. The lines indicate the TDHFB trajectories fulfilling the EBK quantization condition of Eq. (3.7).

A. Stationary-phase approximation to the path integral

Starting an arbitrary state $|\psi(0)\rangle$ at time $t = 0$, the time-dependent full quantum state can be written in the path integral form,

$$\begin{aligned} |\psi(t)\rangle &= e^{-iHt} |\psi(0)\rangle \\ &= \int d\mu(Z'') |Z''\rangle \int d\mu(Z') \\ &\quad \times \int_{Z(0)=Z'}^{Z(t)=Z''} \mathcal{D}\mu[Z(\tau)] e^{iS[Z(\tau)]} \psi(Z'), \end{aligned} \quad (3.1)$$

where $\psi(Z) \equiv \langle Z|\psi(0)\rangle$ and the invariant measure $d\mu(Z)$ is defined by the unity condition,

$$\int d\mu(Z) |Z\rangle \langle Z| = 1. \quad (3.2)$$

In Eq. (3.1), $S[Z(\tau)]$ is the action (2.7) along a given path $Z(\tau)$ with the initial coherent state $|Z(0)\rangle = |Z'\rangle$ and the final state $|Z(t)\rangle = |Z''\rangle$, then, the integration $\int \mathcal{D}\mu[Z(\tau)]$ is performed over all possible paths $|Z(\tau)\rangle$ between them. Among all trajectories in the path integral, the lowest stationary-phase approximation selects the TDHFB (classical) trajectories¹.

$$|\psi(t)\rangle \approx \int d\mu(Z') |Z'_{\text{cl}}(t)\rangle e^{iS_{\text{cl}}(Z'_{\text{cl}}(t), Z')} \psi(Z'), \quad (3.3)$$

where the TDHFB trajectory starting from $|Z'\rangle$ ends at $|Z'_{\text{cl}}(t)\rangle$ at time t . The action $S_{\text{cl}}(Z_f, Z_i)$ is calculated along this classical trajectory connecting $Z_i = Z'_{\text{cl}}(0) = Z'$ and $Z_f = Z'_{\text{cl}}(t)$.

$$\begin{aligned} S_{\text{cl}}(Z'_{\text{cl}}(t), Z') &\equiv \int_0^t \langle Z_{\text{cl}}(t) | i \frac{\partial}{\partial t} - H | Z_{\text{cl}}(t) \rangle dt \\ &= \mathcal{T}[Z_{\text{cl}}] - \mathcal{H}(Z', Z'^*), \end{aligned} \quad (3.4)$$

¹In this formulation, the stationary-phase approximation agrees with the TDHF(B) trajectories, while that to the auxiliary-field path integral of Refs. [28,29] leads to the TDH(B) without the Fock potentials.

with

$$\begin{aligned} \mathcal{T}[Z_{\text{cl}}] &\equiv \int_0^t \langle Z_{\text{cl}}(t) | i \frac{\partial}{\partial t} | Z_{\text{cl}}(t) \rangle dt \\ &= \int_{Z'}^{Z'_{\text{cl}}(t)} \frac{i}{2} \sum_l \frac{\Omega_l}{1 + |Z_l|^2} (Z_l^* dZ_l - Z_l dZ_l^*). \end{aligned} \quad (3.5)$$

In the last equation of Eq. (3.4), we used the fact that the TDHFB trajectory conserves the energy, $\mathcal{H}(Z_{\text{cl}}(t), Z_{\text{cl}}^*(t)) = \mathcal{H}(Z', Z'^*)$.

The energy eigenstates correspond to stationary states, $\langle Z|\psi(t)\rangle \propto \langle Z|\psi(0)\rangle = \psi(Z)$, which can be constructed by superposing the coherent states along a periodic TDHFB trajectory $Z_{\text{cl}}^{(k)}$ as [31,32,39]

$$|\psi_k\rangle = \oint d\mu(Z_{\text{cl}}^{(k)}) |Z_{\text{cl}}^{(k)}\rangle e^{i\mathcal{T}[Z_{\text{cl}}^{(k)}]}. \quad (3.6)$$

The single valuedness of the wave function leads to the quantization condition (k , integer):

$$\begin{aligned} \mathcal{T}_{\circ}[Z_{\text{cl}}^{(k)}] &= \oint \frac{i}{2} \sum_l \frac{\Omega_l}{1 + |Z_l^{(k)}|^2} \\ &\quad \times (Z_l^{(k)*} dZ_l^{(k)} - Z_l^{(k)} dZ_l^{(k)*}) \\ &= 2k\pi. \end{aligned} \quad (3.7)$$

The state evolves in time as $|\psi_k(t)\rangle = |\psi_k\rangle e^{-iE_k t}$, with the energy of the k th periodic trajectory, $E_k = \mathcal{H}(Z_{\text{cl}}^{(k)}, Z_{\text{cl}}^{(k)*})$.

Finding TDHFB trajectories satisfying the quantization condition (3.7) is an extremely difficult task in general. It is better founded and more practical if the classical system is completely integrable. In integrable systems, M complex variables $Z(t)$ can be transformed into the action-angle variables;

$$\begin{aligned} Z(t) &= \{Z_l(t); l = 1, \dots, M\} \\ &\rightarrow \{E; v_1, \dots, v_{M-1}; \theta_1(t), \dots, \theta_M(t)\}, \end{aligned} \quad (3.8)$$

where the variables E and v define an invariant torus, while $\theta(t)$ parametrize the coordinates on the torus. The integration path of Eq. (3.7) is now taken as a topologically independent closed path on the torus, namely the EBK quantization condition. There are M independent closed paths and M quantum

numbers, $k = (k_1, \dots, k_M)$, to specify the stationary energy eigenstate. These are associated with M invariant variables, $\{E_k; v_1^{(k)}, \dots, v_{M-1}^{(k)}\}$. Using the invariant measure,

$$d\mu(Z) = \rho(E, v, \theta) dE dv_1 \dots dv_{M-1} d\theta_1 \dots d\theta_M, \quad (3.9)$$

the k th semiclassical wave function can be calculated as

$$|\psi_k\rangle \propto \oint d\theta_1 \dots \oint d\theta_M \rho(E_k, v^{(k)}, \theta) |E_k, v^{(k)}, \theta\rangle e^{i\mathcal{T}[E_k, v^{(k)}, \theta]}. \quad (3.10)$$

Here, we omit the integration with respect to the invariant variables, E and v .

We apply the semiclassical approach to the two-level pairing model. The invariant measure $\mathcal{U}(2) \otimes \mathcal{U}(2)$ is

$$d\mu(Z) = \prod_l \frac{\Omega_l + 2}{2\pi} (1 + |Z_l|^2)^{-2} d\text{Re} Z_l d\text{Im} Z_l \quad (3.11)$$

$$= \prod_l \frac{\Omega_l + 2}{8\pi} \sin \theta_l d\theta_l d\phi_l \quad (3.12)$$

$$= \left(\prod_{l=1,2} \frac{1 + 2\Omega_l^{-1}}{4\pi} \right) d\Phi dJ d\phi dj. \quad (3.13)$$

In the last equation, we transform the canonical coordinates by Eqs. (2.11) and (2.12). Because the particle number $J = N/2$ and the total energy E are invariant, the two-level pairing model is integrable. Thus, we can construct the semiclassical wave function using Eq. (3.10). The action integral is given by

$$\begin{aligned} \mathcal{T}_k(\Phi, \phi; J) &= J\Phi + \int_{-\pi}^{\phi} j' d\phi' = \frac{N}{2}\Phi + \int_0^t j(t') \frac{d\phi}{dt'} dt' \\ &\equiv \mathcal{T}_{N, E_k}(\Phi, t), \end{aligned} \quad (3.14)$$

where the integration $\int j d\phi$ is performed on the k th closed trajectory of Eq. (3.7), and the variables (ϕ, j) are transformed into (t, E) . The semiclassical wave function fulfilling the EBK quantization condition becomes

$$|\psi_k^N\rangle \propto \oint d\Phi \oint d\phi dt e^{i\mathcal{T}_{N, E_k}(\Phi, t)} |\Phi, t\rangle_{N, E_k} \quad (3.15)$$

$$\propto \sum_{m=0}^J C_m^{(E_k, J)} |S_1, -S_1 + m; S_2, -S_2 + (J - m)\rangle, \quad (3.16)$$

with $S_l = \Omega_l/2$, $J = N/2$, and the coefficients,

$$\begin{aligned} C_m^{(E_k, J)} &= \binom{J}{m} \int_0^T dt \\ &\times \exp \left(i \int_0^t j(t') \dot{\phi}(t') dt' - i(J/2 - m)\phi(t) \right) \\ &\times A(q_1, S_1, m) A(q_2, S_2, J - m), \quad (3.17) \\ A(q, S, m) &= \left(\frac{1-q}{2} \right)^{m/2} \left(\frac{1+q}{2} \right)^{S-m/2} \sqrt{\frac{(2S)! m!}{(2S-m)!}}, \end{aligned}$$

where T is the period of the closed trajectory. The TDHFB-requantized wave functions (3.16) are eigenstates of the total

particle number. This is because of the integration over the global gauge angle Φ , which makes the particle number projection not only for the ground state but also for excited states. See appendix for detailed derivation of Eq. (3.16).

Because we obtain the microscopic wave function of every eigenstate, the expectation values and the transition matrix elements for any operator can be calculated in a straightforward manner. In Sec. IV, we show those of the pair-addition operator S^+ which characterize properties of the pair condensates.

Before ending this section, let us note the periodic conditions of the coordinates and the quantization condition. Because the two original variables, (χ_1, χ_2) , are independent periodic variables of the period 2π , in addition to the trivial periodicity of 2π for Φ and of 4π for ϕ , we have periodic conditions for $(\Phi, \phi) \rightarrow (\Phi \pm \pi, \phi \pm 2\pi)$ and $(\Phi, \phi) \rightarrow (\Phi \pm \pi, \phi \mp 2\pi)$. The former (latter) corresponds to $\chi_2 \pm 2\pi$ ($\chi_1 \pm 2\pi$) with χ_1 (χ_2) being fixed. For open TDHFB trajectory (e.g., Fig. 1), the quantization condition becomes

$$\begin{aligned} \mathcal{T}_{N, E_k}(\Phi \pm \pi, \phi \pm 2\pi; J) &= \mathcal{T}_{N, E_k}(\Phi, \phi; J) + 2m\pi \\ \Leftrightarrow \quad \pm \frac{N}{2}\pi + \int_{-\pi}^{\pi} j d\phi &= 2m\pi. \end{aligned} \quad (3.18)$$

This leads to the following:

$$\int_{-\pi}^{\pi} j d\phi = \begin{cases} 2k\pi & \text{for } N = 4n, \\ (2k+1)\pi & \text{for } N = 4n+2, \end{cases} \quad (3.19)$$

where m, k , and $n > 0$ are integer numbers.

B. Canonical quantization

The most common approach to the quantization of the nuclear collective model is the canonical quantization [1]. In the pairing collective model, the canonical quantization was adopted in previous studies [34–36]. Assuming magnitude and phase of the pairing gap as collective coordinates, a collective Hamiltonian was constructed in the second order in momenta. Then, the Hamiltonian was quantized by the canonical quantization with Pauli's prescription. In this section, we apply a similar quantization method to the TDHFB Hamiltonian (2.15). The main difference is that the collective canonical variables are not assumed in the present case, but are obtained from the TDHFB dynamics itself.

It is not straightforward to apply Pauli's prescription to the present case, because the TDHFB Hamiltonian (2.15) is not limited to the second order in momenta. In the present study, we adopt a simple symmetrized ordering, as

$$\begin{aligned} H(\hat{\phi}, \hat{j}, \hat{J}) &= \sum_{l=1,2} \Omega_l \epsilon_l (1 - q_l) \\ &- \frac{g}{4} \sum_{l=1,2} \Omega_l (\Omega_l (1 - q_l^2) + (1 - q_l)^2) \\ &- \frac{g}{4} \Omega_1 \Omega_2 \left\{ \sqrt{(1 - q_1^2)(1 - q_2^2)} \cos \hat{\phi} \right. \\ &\left. + \cos \hat{\phi} \sqrt{(1 - q_1^2)(1 - q_2^2)} \right\}. \end{aligned} \quad (3.20)$$

As in Eq. (2.16), q_l contain J and j which are replaced by

$$\hat{J} = -i \frac{\partial}{\partial \Phi}, \quad \hat{j} = -i \frac{\partial}{\partial \phi}. \quad (3.21)$$

Because Φ is a cyclic variable, we write the collective wave function $\Psi(\Phi, \phi)$ as eigenstates of the particle number N in a separable form,

$$\Psi_k^{(N)}(\Phi, \phi) = \frac{1}{\sqrt{2\pi}} e^{i \frac{N}{2} \Phi} \psi_k^{(N)}(\phi). \quad (3.22)$$

Then, the problem is reduced to the one-dimensional Schrödinger equation for motion in the relative angle ϕ . The Schrödinger equation,

$$H\left(\phi, -i \frac{d}{d\phi}; \frac{N}{2}\right) \psi_k^{(N)}(\phi) = E_k^{(N)} \psi_k^{(N)}(\phi), \quad (3.23)$$

The wave function should have a periodic property with respect to the variable ϕ ; $\psi_k(\phi) = \psi_k(\phi + 4\pi)$. For the adopted simple ordering of Eq. (3.20), it is convenient to use the eigenstates of \hat{j} as the basis to diagonalize the Hamiltonian. They are

$$\chi_j(\phi) = \frac{1}{\sqrt{4\pi}} e^{i\phi j}, \quad \text{with } j: \text{integer or half integer.} \quad (3.24)$$

Because the Hamiltonian (3.20) contains only terms linearly proportional to $e^{\pm i\phi}$, the basis states χ_j with half-integer difference in j are not coupled with each other. Thus, the eigenstates of Eq. (3.23) can be expanded as

$$\psi_k^{(N)}(\phi) = \sum_{j=j_{\min}, j_{\min}+1, \dots}^{j_{\max}} c_{k,j}^{(N)} \chi_j^{(N)}(\phi). \quad (3.25)$$

According to the relation $j = (n_2 - n_1)/4 = (N - 2n_1)/4$ in Eq. (2.14), we adopt the (half-)integer values of j for $N = 4n$ ($N = 4n + 2$) with integer n . This is consistent with the quantization condition (3.19). The coupling term with different j in Eq. (3.20) vanishes for $n_l = 0$ and $n_l = 2Q_l$, which restricts values of j in a finite range of $j_{\min} \leq j \leq j_{\max}$.

To estimate the two-particle transfer matrix elements, we construct the corresponding operators as follows. The classical form of matrix elements are obtained as

$$S^+(\Phi, J; \phi, j) = \langle Z | \hat{S}^+ | Z \rangle = \frac{1}{2} (\Omega_1 \sqrt{1 - q_1^2} e^{-i\phi/2} + \Omega_2 \sqrt{1 - q_2^2} e^{i\phi/2}) e^{i\Phi}, \quad (3.26)$$

$$S^-(\Phi, J; \phi, j) = \langle Z | \hat{S}^- | Z \rangle = \frac{1}{2} (\Omega_1 \sqrt{1 - q_1^2} e^{i\phi/2} + \Omega_2 \sqrt{1 - q_2^2} e^{-i\phi/2}) e^{-i\Phi}. \quad (3.27)$$

Again, we adopt a simple symmetrized ordering for the quantization:

$$\begin{aligned} S^\pm(\hat{\Phi}, \hat{J}; \hat{\phi}, \hat{j}) &= \frac{1}{4} (\Omega_1 \sqrt{1 - q_1^2} e^{\mp i\hat{\phi}/2} + \Omega_2 \sqrt{1 - q_2^2} e^{\pm i\hat{\phi}/2}) \\ &\times e^{\pm i\hat{\Phi}} + \frac{1}{4} e^{\pm i\hat{\Phi}} (e^{\mp i\hat{\phi}/2} \Omega_1 \sqrt{1 - q_1^2} \\ &+ e^{\pm i\hat{\phi}/2} \Omega_2 \sqrt{1 - q_2^2}). \end{aligned} \quad (3.28)$$

The exponential factors $e^{\pm i\Phi}$ change the total particle number $N \rightarrow N \pm 2$, while $e^{\pm i\phi/2}$ change the relative numbers, $n_2 - n_1 \rightarrow n_2 - n_1 \pm 2$. Using these operators, the pair-addition transition strengths are calculated as

$$\begin{aligned} B(P_{\text{ad}}; k \rightarrow k') &= |\langle N', k' | S^+(\hat{\Phi}, \hat{J}; \hat{\phi}, \hat{j}) | N, k \rangle|^2 \\ &= \left| \int_0^{2\pi} d\Phi \int_{-2\pi}^{2\pi} d\phi \psi_{k'}^{(N')*}(\phi) e^{-i \frac{N'}{2} \Phi} S^+ \right. \\ &\quad \left. \times (\hat{\Phi}, \hat{J}; \hat{\phi}, \hat{j}) \psi_k^{(N)}(\phi) e^{i \frac{N}{2} \Phi} \right|^2, \end{aligned} \quad (3.29)$$

which automatically vanishes for $N' \neq N + 2$.

C. Fourier decomposition of time-dependent matrix elements

The requantization and calculation of the matrix elements also can be performed using the time-dependent solutions of the TDHFB. It was proposed and applied to the two-level pairing model [38], which we recapitulate in this section.

The TDHFB provides a time-dependent solution $Z(t)$ starting from a given initial state $Z(0)$. The energy eigenvalues and the corresponding closed trajectories are determined from the EBK quantization condition (3.7). The pair-transfer matrix elements are evaluated as the Fourier components of the time-dependent mean values $S^\pm(t) = S^\pm(Z(t))$, Eqs. (3.26) and (3.27). Because the global gauge angle Φ is a cyclic variable, the motion in the relative gauge angle ϕ is independent from Φ . Thus, we calculate the time evolution of $\phi(t)$, and find the period of the k th closed trajectory T satisfying Eq. (3.7). Then, the Fourier component,

$$\tilde{S}^\pm(E_k; \omega) = \frac{1}{T} \int_0^T dt e^{i\omega t} S^\pm(t), \quad (3.30)$$

corresponds to the pair-transfer matrix element from the state k to k' when $\omega = 2\pi(k' - k)/T$. The pair-addition transition strengths are calculated as

$$B(P_{\text{ad}}; k \rightarrow k') = \left| \tilde{S}^+ \left(E_k; \frac{2\pi}{T} \Delta k \right) \right|^2, \quad (3.31)$$

with $\Delta k = k' - k$. In this approach, the transition between the ground states of neighboring nuclei ($N \rightarrow N + 2$) corresponds to the stationary component ($k = 0$ and $\Delta k = 0$), namely the expectation value in the BCS approximation.

The derivation of Eq. (3.30) is based on the wave packet in the classical limit [43]. The TDHFB state is assumed to be a superposition of eigenstates $|\phi_k^N\rangle$ in a narrow range of energy $E_{k_0} - \Delta E < E_k < E_{k_0} + \Delta E$,

$$|Z(t)\rangle = \sum_N \sum_k c_k^N |\phi_k^N\rangle e^{-iE_k t}, \quad (3.32)$$

where the eigenenergies are evenly spaced and the coefficients c_k^N slowly vary with respect to k and N . The expectation value of S^\pm is

$$S^\pm(t) = \sum_N \sum_{k, k'} c_{k'}^{N+2*} c_k^N \langle \phi_{k'}^{N+2} | S^\pm | \phi_k^N \rangle e^{i(E_{k'} - E_k)t}. \quad (3.33)$$

The matrix element $\langle \phi_{k'}^{N+2} | S^\pm | \phi_k^N \rangle$ quickly disappears as $|k' - k|$ increases, while it stays almost constant for the

small change of k and N with $|k' - k|$ being fixed. Thus, we may approximate $c_{k'}^{N+2} \approx c_k^N$, $E_{k'} - E_k \approx \omega_0 \Delta k$, and that $\langle \phi_{k'}^{N+2} | S^\pm | \phi_k^N \rangle \approx \langle \phi_{k_0+\Delta k}^{N+2} | S^\pm | \phi_{k_0}^N \rangle$,

$$\begin{aligned} S^\pm(t) &\approx \sum_N \sum_k |c_k^N|^2 \sum_{\Delta k} \langle \phi_{k_0+\Delta k}^{N+2} | S^\pm | \phi_{k_0}^N \rangle e^{i\omega_0 \Delta k t} \\ &= \sum_{\Delta k} \langle \phi_{k_0+\Delta k}^{N+2} | S^\pm | \phi_{k_0}^N \rangle e^{i\omega_0 \Delta k t}, \end{aligned} \quad (3.34)$$

where k_0 is a representative index value of the superposition in Eq. (3.32). From this classical wave packet approximation, we obtain Eq. (3.30). It is not trivial to justify the approximation for small values of Ω and for transitions around the ground state.

IV. RESULTS

In this section, we study the seniority-zero states ($\nu_1 = \nu_2 = 0$) in the two-level system with equal degeneracy, $\Omega_1 = \Omega_2 = \Omega$. Because all the properties are scaled with the ratio $g/\Delta\epsilon$, where $\Delta\epsilon$ is the level spacing $\Delta\epsilon = \epsilon_2 - \epsilon_1$, we define a dimensionless parameter to control the strength of the pairing correlation,

$$x = 2\Omega \frac{g}{\Delta\epsilon}. \quad (4.1)$$

For sub-shell-closed systems with the $N = 2\Omega$ system, the transition from normal ($x < 1$) to superfluid ($x > 1$) states takes place at $x = 1$.²

We apply the requantization methods in Sec. III. In the following, the stationary-phase approximation to the path integral in Sec. III A is denoted as “SPA,” the Fourier decomposition method (Sec. III C) as “FD,” and the canonical quantization with periodic boundary condition (Sec. III B) as “CQ.” Note that the SPA and the FD produce the same eigenenergies which are based on the EBK quantization rule.

A. Large- Ω cases

In the limit of $\Omega \rightarrow \infty$, we expect that the classical approximation becomes exact. Here, we adopt $\Omega = 50$ with $N = 100$ (closed-shell configuration) and $N = 50$ (mid-shell configuration).

Calculated excitation energies are shown in Fig. 2. The results of SPA/FD and CQ are compared with the exact values. At the weak pairing limit of $x \rightarrow 0$, the excitation energies are multiples of $2\Delta\epsilon$, which correspond to pure $2n$ -particle- $2n$ -hole excitations. Both the weak and the strong pairing limits are nicely reproduced by all the calculations, while the CQ method produces excitation energies slightly lower than the exact values in an intermediate region around $x = 1$. It is somewhat surprising to see that the deviation is larger for the case of the mid-shell configuration ($N = 50$) than the closed shell ($N = 100$).

The deviation in the CQ method is mainly from the zero-point energy in the ground state. Because we solve the




FIG. 2. Excitation energies of $|0_2^+\rangle$, $|0_3^+\rangle$, and $|0_4^+\rangle$ for $\Omega = 50$ systems with (a) $N = 50$ and (b) $N = 100$ as functions of the dimensionless parameter x of Eq. (4.1).

collective Schrödinger equation (3.23) with the quantized Hamiltonian of Eq. (3.20), the zero-point energy $\Delta E > 0$ is inevitable in the CQ method. The ΔE is associated with the degree of localization of the wave function. Thus, the magnitude of ΔE for “bound” states is different from that for “unbound” states. See Fig. 1. In the strong pairing limit, the potential minimum is deep enough to bound both ground and excited states. Conversely, all the states are unbound in the weak limit. In both limits, ΔE for ground and excited states are similar, and they are canceled for the excitation energy. However, near $x = 1$, the ground state is bound, while the excited states are unbound. In this case, ΔE is larger in the ground state than in the excited states, which makes the excitation energy smaller. This also explains the difference between the mid-shell and closed-shell configurations. In the closed shell, all the states are unbound for $x < 1$, while, in the mid-shell, there is a region in $x < 1$ where the ground state is bound but the excited state is unbound.

The obtained wave functions in the SPA and the CQ can be decomposed in the $2n$ -particle- $2n$ -hole components in Fig. 3. In the SPA, it is done as Eq. (3.16) and the normalized squared coefficients $|C_m^{(E_k, J)}|^2$ are plotted in Fig. 3. For the CQ, $|c_{k,j}^{(N)}|^2$ in Eq. (3.25) are shown. Here, m and j are related to each other, $2j = J - 2m$. They show excellent agreement with the exact

²Strictly speaking, the phase transition takes place at $x = 2\Omega/(2\Omega - 1)$.




FIG. 3. Occupation probability in excited 0^+ states as a function of j for $\Omega = 50$ systems with (a) $N = 50$ and (b) $N = 100$. The upper and lower panels display the results for $x = 0.5$ and $x = 2$, respectively. The three vertical bars at each j from the left to the right represent the squared components of the wave functions from exact, SPA, and CQ calculations, respectively. The left end of the horizontal axis at $j = j_{\min}$ corresponds to a component with $(n_1, n_2) = (N, 0)$. The next at $j = j_{\min} + 1$ corresponds to the one with $(n_1, n_2) = (N - 2, 2)$, and so on.

results, not only for the ground state but for excited states. We find the SPA is even more precise than the CQ.

Next, let us discuss the transition matrix elements. In this paper, we discuss only $k = 0$ (ground state) and $k = 1$ (first excited $\nu = 0$ state). The FD calculation is based on the time evolution of the expectation value $S^+(t)$ with fixed (J, Φ) in Eq. (3.30). For $(N, k) \rightarrow (N + 2, k')$ transitions, we basically adopt the trajectories for the initial state, namely, the one with $J = N/2$ satisfying the k th EBK quantization condition. The $k \rightarrow k$ ($\Delta k = 0$) transitions correspond to the intraband transitions of the pair-rotational band, when the state is deformed in the gauge space (pair deformation). For the ground-state band ($k = 0$), this is nothing but the expectation value at the BCS wave function, with the constant value of S^+ . Because the constant S^+ provides only $\Delta k = 0$ intraband transitions, for the interband transition of $(N, k = 0) \rightarrow (N + 2, k = 1)$ transitions, the trajectory satisfying the EBK condition of $k = 1$ is used to perform the Fourier decomposition (3.30) of $\omega = 2\pi/T$.

The calculated pair-addition strengths $B(P_{\text{ad}})$ are shown in Fig. 4 for $N = 48 \rightarrow 50$, and in Fig. 5 for $N = 98 \rightarrow 100$. Near the closed-shell configuration ($N = 98 \rightarrow 100$), the pair-addition strengths for the intraband transitions ($\Delta k = 0$)

drastically increase around $x = 1$. This reflects a character change from the pair vibration ($x \lesssim 1$) to the pair rotation ($x \gtrsim 1$). The $B(P_{\text{ad}}; k \rightarrow k)$ in the pair-rotational transitions are about 20 times larger than those in the vibrational transitions. The interband $B(P_{\text{ad}}; 0 \rightarrow 1)$ are similar to the $B(P_{\text{ad}}; 0 \rightarrow 0)$ in the vibrational region ($x \lesssim 1$), because they both change the number of pair-phonon quanta by one unit. In contrast, $B(P_{\text{ad}}; 1 \rightarrow 0)$, which change the phonon quanta by three, are almost zero. In the pair-rotational region ($x \gg 1$), $B(P_{\text{ad}}; 1 \rightarrow 0)$, and $B(P_{\text{ad}}; 0 \rightarrow 1)$ are roughly identical. This is because both $B(P_{\text{ad}}; 1 \rightarrow 0)$ and $B(P_{\text{ad}}; 0 \rightarrow 1)$ correspond to one-phonon excitation in “deformed” cases ($x \gg 1$).

In the mid-shell region ($N = 48 \rightarrow 50$), the intraband $B(P_{\text{ad}}; k \rightarrow k)$ smoothly increases as x increases. Their values are larger than the interband strengths by about one (two) order of magnitude at $x \sim 0$ ($x \sim 2.5$), indicating the pair-rotational character. The interband $B(P_{\text{ad}}; 0 \rightarrow 1)$ shows a gradual decrease as a function of x , while $B(P_{\text{ad}}; 1 \rightarrow 0)$ is negligibly small, even at $x \gg 1$. This presents a prominent difference from the closed-shell case.

All the features of the pair-transfer strengths are nicely reproduced in the SPA method, for both the closed- and mid-shell configurations. The CQ method qualitatively agrees with




FIG. 4. The strength of pair-addition transition $B(P_{ad}; k \rightarrow k')$ for $\Omega = 50$ systems from $N = 48 \rightarrow 50$. (Left panels) Results of the CQ method; middle panels, FD; right panels, SPA. Dashed lines represent exact calculation. Upper panels show the intraband transitions of $|0_1^+\rangle \rightarrow |0_1^+\rangle$ and $|0_2^+\rangle \rightarrow |0_2^+\rangle$, while lower panels show the interband transition of $|0_1^+\rangle \rightarrow |0_2^+\rangle$ and $|0_2^+\rangle \rightarrow |0_1^+\rangle$.

the exact calculation. For instance, the order-of-magnitude difference between intraband and interband transitions. However, the precision of the CQ method is not so good, especially around $x = 1$. The FD method properly describes the main features in the superfluid phase, while it fails for the normal phase ($x \lesssim 1$). In the mid-shell configuration, the ground state is always in the superfluid phase at $x > 0$, while the $k = 1$ excited state corresponds to the open (closed) trajectory at $0 < x \lesssim 1$ ($x \gtrsim 1$). For the open trajectory, the FD produces wrong values. However, somewhat surprisingly, the SPA, which uses these open trajectories for the construction of wave functions, reproduces main features of the exact results.

B. Small- Ω cases

Next, we discuss systems with smaller degeneracy $\Omega = 8$. Again, we study systems near the closed-shell and the mid-shell configurations.

1. Mid-shell configuration

The calculated excitation energies are shown in Fig. 6 for the $N = 8$ case. The SPA/FD reproduces the exact calculation in the entire region of x , not only for the lowest but also for higher excited states. The CQ reproduces the exact result in a weak pairing region ($x \lesssim 1$), while it underestimates the excitation energies at $x \gtrsim 1$. Analogous to the case of $\Omega = 50$,




FIG. 5. The same as Fig. 4 but for $N = 98 \rightarrow 100$.




FIG. 6. Excitation energies of $|0_2^+)$, $|0_3^+)$, and $|0_4^+)$ for $\Omega = N = 8$ systems as functions of x .





FIG. 7. Occupation probability in excited 0^+ states as a function of j for $\Omega = N = 8$ systems. (a) and (b) Results for $x = 0.5$ and $x = 2$, respectively. See also the caption for Fig. 3.

(SPA/FD and CQ). The behaviors of the lowest excitation agree with the exact calculations, while the CQ method substantially underestimates those for higher states. This is again from the difference in the zero-point energy in the ground and the excited states. In the CQ calculation, the first excited state is bound at $x \gtrsim 2$, but the second excited state is unbound for $x \lesssim 3.2$.

Near the transition point from the open to closed trajectories, the wave functions calculated with the SPA and CQ methods somewhat differ from the exact ones. In Fig. 10, the wave functions at $x = 0.5$ and 2 are presented. They agree with exact calculation at $x = 0.5$. In contrast, we find some deviations for the first excited state ($k = 1$) at $x = 2$. This is because the $k = 1$ trajectory corresponding to the first excited state changes its character from open to closed at $x \approx 1.8$. Therefore, the first excited wave function is difficult to reproduce in the SPA, although the wave functions for the ground and higher excited states show reasonable agreement. The similar disagreement is observed for the ground state near $x = 1$.

Singular behaviors near the transition points also can be observed in the pair-addition transition strengths ($N = 14 \rightarrow 16$) shown in Fig. 11. At $x = 1$, the intraband $B(P_{ad}; 0 \rightarrow 0)$ shows

this is mainly from the effect of the zero-point energy ΔE . The ground-state energy in the CQ calculation is bound at $x \gtrsim 1$. However, because of the weak collectivity with $N = 8$, the first excited state stays unbound even at the maximum x in Fig. 6. Therefore, the energy shift $\Delta E > 0$ is larger in the ground state, which makes the excitation energy smaller.

The wave functions are plotted in Fig. 7. At the weak pairing case of $x = 0.5$, both the SPA and the CQ reproduce the exact result. At $x = 2$, the squared coefficients of the ground state has an asymmetric shape peaked at the lowest j , which suggests that the state is not deeply bound in the potential. It is in contrast to the symmetric shape in Fig. 3. The wave functions obtained by the CQ method has noticeable deviation from the exact results. On the other hand, the SPA wave functions are almost identical to the exact ones.


The pair-addition transition strengths from $N = 6$ to $N = 8$ are shown in Fig. 8. The intraband $k \rightarrow k$ transitions increase and the interband $k = 0 \rightarrow 1$ transitions decrease as functions of x . Their relative difference becomes more than one order of magnitude at $x \gtrsim 2$. Thus, even at relatively small Ω and N , the intraband transitions in the pair rotation is qualitatively different from the interband transitions.

We find the excellent agreement between the SPA and the exact calculations. The first excited state corresponds to the open trajectory which turns out to almost perfectly reproduce the exact wave function. In contrast, this open trajectory produces results far from the exact one in the FD method. It produces almost vanishing the intraband $B(P_{ad}; 1 \rightarrow 1)$. The $B(P_{ad}; 0 \rightarrow 0)$ shows a qualitative agreement for its behavior, but is significantly underestimated. The CQ method also underestimates the intraband transitions.

For the mid-shell configurations, the SPA is dominantly superior to the CQ and the FD methods.

2. Closed-shell configuration

In the closed shell with $N = 16$, the minimum-energy trajectory changes at $x = 1$ from $j = -4$ (normal phase) to the BCS minimum $j > -4$ and $\phi = 0$ (superfluid phase). At the transitional point ($x = 1$), the harmonic approximation is known to collapse, namely to produce zero excitation energy. In Fig. 9, this collapsing is avoided in all the calculations


FIG. 8. The same as Fig. 4 but for $N = 6 \rightarrow 8$ with $\Omega = 8$.

a kink in the SPA, and $B(P_{ad}; 1 \rightarrow 1)$ shows another kink at $x \approx 1.8$. These exactly correspond to the transition points from open to closed trajectories. Nevertheless, the overall behaviors are well reproduced and the values at the weak and strong pairing limit are reasonably reproduced in the SPA. The CQ calculation also shows smoothed kinklike behaviors near the transition points. However, it underestimates the intraband $B(P_{ad}; k \rightarrow k)$. The FD method does not have a kink for $B(P_{ad}; 0 \rightarrow 0)$, because $S^+(t)$ is calculated for an $N = 14$ system. Both intraband and interband transitions in the FD calculations reasonably agree with the exact results at $x \gtrsim 1.8$. The $k = 1$ state is not properly reproduced at $x \lesssim 1.8$ with the open trajectory.


For the closed-shell configurations, the SPA and the FD methods provide reasonable description for the pair-transfer transition strengths.


C. Collective model treatment

The collective model was proposed and utilized for the nuclear pairing dynamics [34–36]. For those studies, the

FIG. 9. The same as Fig. 6 but for $N = 2\Omega = 16$.

pairing gap parameter (or equivalent quantities) is assumed to be the collective coordinates. This is analogous to the five-dimensional (5D) collective (Bohr) model, in which the

FIG. 10. The same as Fig. 7 but for $N = 2\Omega = 16$.

FIG. 11. The same as Fig. 4 but for $N = 14 \rightarrow 16$ with $\Omega = 8$.

collective coordinates are assumed to be the quadrupole deformation parameters $\alpha_{2\mu}$. The 5D collective model was extensively applied to analysis on numerous experimental data. On the contrary, there have been very few applications of the pairing collective model in comparison with experimental data. In this section, we examine the validity of the collective treatment of the pairing.

Although the global gauge angle Φ is arbitrary, the deformation parameter $\alpha \equiv \langle \hat{S}^- \rangle$ in the gauge space is usually taken as a real value ($\Phi = 0$). The energy minimization with a fixed value of real α always leads to $\phi = 0$.


$$\alpha(j, J) = \langle Z_0 | \hat{S}^- | Z_0 \rangle = \frac{\Omega}{2} (\sqrt{1 - q_1^2} + \sqrt{1 - q_2^2}). \quad (4.2)$$

The parameter α is equivalent to the pairing gap Δ , because the relation, $\Delta = G\alpha$, guarantees one-to-one correspondence between α and Δ . In Sec. III B, we treat ϕ as a collective coordinate and j as its conjugate momentum. The collective model treatment is based on the opposite choice, j as a coordinate and ϕ as a momentum.

The problem is that there is no one-to-one correspondence between j and α . The relation between j and α are shown by dashed lines in Fig. 12 for $\Omega = 8$ mid-shell (a) and closed-shell (b) configurations. The deformation parameter α is largest at $j = 0$ (equal filling in both levels), and smallest at the end points of j . The constrained minimization with respect to α cannot produce the states corresponding to $j > 0$. Apparently, we cannot map the entire region of j to α .

The collective model treatment requires the collective wave functions to be well localized in the $j < 0$ region. The potential energy, $E(j) = \mathcal{H}(\phi = 0, j; J = N/2)$ of Eq. (2.15), is also shown in Fig. 12. The restriction becomes more serious for the stronger pairing cases. For instance, the potential with $x = 3.2$ in Fig. 12(a) has only about 1-MeV depth at the minimum point, relative to the value at the boundary point ($j = 0$) corresponding to the maximum value of α (Δ).

To simulate the result of the collective model, we expand the Hamiltonian (2.15) up to the second order in ϕ , then, quantize it by $\hat{\phi} = i\partial/\partial j$ with the ordering given by Pauli's prescription. The range of the coordinate j is restricted to $j_{\min} \leq j \leq 0$

FIG. 12. Energy surfaces as functions of j , Eq. (2.15) with $\phi = 0$, for $x = 0.5, 2$, and 3.2 . Dashed line is the pairing deformation parameter α of Eq. (4.2) as a function of j . (a) Mid-shell ($\Omega = N = 8$) and (b) closed-shell ($2\Omega = N = 16$).




FIG. 13. Potential energy surface as functions of α with $x = 2.4$. (a) $\Omega = N = 50$, (b) $2\Omega = N = 16$. Horizontal lines indicate energy spectra. Black lines are obtained from the potential energy surface, and green lines are obtained with the CQ method (Sec. III B).

(FD) of the time-dependent observables. In this model, because the global gauge angle Φ is a cyclic variable, the TDHFB dynamics can be described by the integrable classical dynamics. After the pair-rotation variables (Φ, J) are separated, the remaining degrees of freedom (ϕ, j) describe the pair-vibrational motion.

In systems with large degeneracy Ω and number of particles N , all the quantization methods reasonably reproduce the results of the exact calculation for excitation spectra. It is more difficult to reproduce the two-particle transfer matrix elements. Nevertheless, for the large Ω and N , we obtain qualitative agreement with the exact results. These are ideal cases, but realistic situations may have smaller Ω and N in the valence space.

In systems with relatively small Ω and N , the agreement is less quantitative for the CQ and the FD, especially for the two-particle transfer matrix elements. In contrast, the SPA keeps its accuracy in the entire range of pairing strengths. One of the reasons of its success is because of the inclusion of the off-diagonal parts of the pair-transfer operator, by the explicit construction of the microscopic wave functions. The CQ and FD calculate the pair-transfer matrix elements using only the diagonal part (expectation value) of the operator \hat{S}^\pm , based on Eqs. (3.26) and (3.27). This is a good approximation when the collectivity is so large that the diagonal parts dominate. However, the pairing collectivity may be too weak to justify this treatment.

We also investigated the conventional treatment of the collective model which assumes that the collective coordinate is the pairing gap parameter. As we mentioned before, the present two-level pairing Hamiltonian has only one pair of collective variables (ϕ, j) , in addition to the pair-rotational variables (Φ, J) . Even for such a simple system, we find that it is difficult to justify the use of Δ as the collective coordinate, especially for relatively small- Ω cases. Basically, there is no one-to-one correspondence between Δ and j . The collective wave functions are not necessarily bound in the region where the variable Δ can represent.

Among the different requantization methods, the SPA is the most accurate tool for description of the pairing large amplitude collective motion in realistic nuclear systems. The weak point of this approach is that it is applicable only to the integrable TDHFB system. To solve this problem, we plan to first extract the integrable collective submanifold in the many-dimensional TDHFB phase space. For this purpose, the adiabatic self-consistent collective coordinate (ASCC) method [8] is a promising tool. The combined study of the ASCC and the SPA for multilevel systems is our next target under progress.

ACKNOWLEDGMENTS

This work is supported in part by Interdisciplinary Computational Science Program in CCS, University of Tsukuba, and by JSPS-NSFC Bilateral Program for Joint Research Project on Nuclear mass and life for unravelling mysteries of the r-process.

V. CONCLUSION

The different methods of the requantization of the TDHFB dynamics was studied for the two-level pairing model; the stationary-phase approximation (SPA) of the path integral, the canonical quantization (CQ), and the Fourier decomposition

APPENDIX: DERIVATION OF SEMICLASSICAL WAVE FUNCTION

We give a derivation of the semiclassical wave function of Eq. (3.16). The explicit form of coherent state is

$$\begin{aligned} |Z\rangle &= \prod_{l=1}^2 (1 + |Z_l|^2)^{-\Omega_l/2} e^{Z_l S_l^+} |0\rangle = \prod_{l=1}^2 \left(1 + \tan^2 \frac{\theta_l}{2}\right)^{-\Omega_l/2} \sum_k \frac{1}{k!} \sum_{m=0}^k \binom{k}{m} \left(\tan \frac{\theta_1}{2} e^{-i\phi_1} S_1^+\right)^m \left(\tan \frac{\theta_2}{2} e^{-i\phi_2} S_2^+\right)^{k-m} |0\rangle \\ &= \prod_{l=1}^2 \left(1 + \tan^2 \frac{\theta_l}{2}\right)^{-\Omega_l/2} \sum_k \frac{1}{k!} \sum_{m=0}^k \binom{k}{m} \tan^m \frac{\theta_1}{2} \tan^{k-m} \frac{\theta_2}{2} e^{-ik\Phi} e^{-i(k/2-m)\phi} (S_1^+)^m (S_2^+)^{k-m} |0\rangle. \end{aligned} \quad (\text{A1})$$

Inserting (A1) into (3.15) under fixed N and E_k , it becomes

$$\begin{aligned} |\psi_k^N\rangle &\propto \oint d\Phi \oint dt e^{i\mathcal{T}_{N,E_k}(\Phi,t)} |\Phi,t\rangle_{N,E_k} \propto \sum_k \frac{1}{k!} \sum_{m=0}^k \binom{k}{m} \int_0^{2\pi} d\Phi e^{i(N/2-k)\Phi} \int_0^T dt e^{i \int \pi(t') \dot{\phi}(t') dt' - i(k/2-m)\phi} \\ &\quad \times \left\{ \prod_{l=1}^2 \left(1 + \tan^2 \frac{\theta_l}{2}\right)^{-\Omega_l/2} \right\} \tan^m \frac{\theta_1}{2} \tan^{k-m} \frac{\theta_2}{2} (S_1^+)^m (S_2^+)^{k-m} |0\rangle \\ &\propto \sum_{m=0}^{N/2} \binom{N/2}{m} \int_0^T dt \exp \left(i \int \pi(t') \dot{\phi}(t') dt' - i(N/4 - m)\phi \right) \\ &\quad \times \left\{ \prod_{l=1}^2 \left(1 + \tan^2 \frac{\theta_l}{2}\right)^{-\Omega_l/2} \right\} \tan^m \frac{\theta_1}{2} \tan^{N/2-m} \frac{\theta_2}{2} (S_1^+)^m (S_2^+)^{N/2-m} |0\rangle. \end{aligned} \quad (\text{A2})$$

We find that the integration over Φ is nothing but the number projection. In SU(2) quasispin representation, the vacuum state is written as $|0\rangle = |S_1, -S_1; S_2, -S_2\rangle$, which leads to

$$(S_1^+)^m (S_2^+)^{N/2-m} |0\rangle = \sqrt{\frac{(2S_1)!m!}{(2S_1-m)!}} \sqrt{\frac{(2S_2)!(N/2-m)!}{[2S_2-(N/2-m)]!}} |S_1, -S_1+m; S_2, -S_2+(N/2-m)\rangle. \quad (\text{A3})$$

For convenience, we define coefficients,

$$A(q, S, m) \equiv \frac{\tan^m \frac{\theta}{2}}{(1 + \tan^2 \frac{\theta}{2})^{-S}} \sqrt{\frac{(2S)!m!}{(2S-m)!}} = \left(\frac{1-q}{2}\right)^{m/2} \left(\frac{1+q}{2}\right)^{S-m/2} \sqrt{\frac{(2S)!m!}{(2S-m)!}}, \quad (\text{A4})$$

where $q = \cos \theta$. Inserting Eq. (A4) into Eq. (A2) with $N/2 = J$, we reach Eq. (3.16).

-
- [1] A. Bohr and B. R. Mottelson, *Nuclear Structure, Vol. II* (W. A. Benjamin, New York, 1975).
- [2] P. Ring and P. Schuck, *The Nuclear Many-body Problems*, Texts and Monographs in Physics (Springer-Verlag, New York, 1980).
- [3] D. Brink and R. A. Broglia, *Nuclear Superfluidity, Pairing in Finite Systems* (Cambridge University Press, Cambridge, 2005).
- [4] P. E. Garrett, *J. Phys. G* **27**, R1 (2001).
- [5] K. Heyde and J. L. Wood, *Rev. Mod. Phys.* **83**, 1467 (2011).
- [6] M. Bender, P. H. Heenen, and P.-G. Reinhard, *Rev. Mod. Phys.* **75**, 121 (2003).
- [7] J.-P. Blaizot and G. Ripka, *Quantum Theory of Finite Systems* (MIT Press, Cambridge, 1986).
- [8] T. Nakatsukasa, K. Matsuyanagi, M. Matsuo, and K. Yabana, *Rev. Mod. Phys.* **88**, 045004 (2016).
- [9] S. Ebata, T. Nakatsukasa, T. Inakura, K. Yoshida, Y. Hashimoto, and K. Yabana, *Phys. Rev. C* **82**, 034306 (2010).
- [10] I. Stetcu, A. Bulgac, P. Magierski, and K. J. Roche, *Phys. Rev. C* **84**, 051309 (2011).
- [11] Y. Hashimoto, *Eur. Phys. J. A* **48**, 55 (2012).
- [12] G. Scamps and D. Lacroix, *Phys. Rev. C* **88**, 044310 (2013).
- [13] S. Ebata, T. Nakatsukasa, and T. Inakura, *Phys. Rev. C* **90**, 024303 (2014).
- [14] G. Scamps and D. Lacroix, *Phys. Rev. C* **87**, 014605 (2013).
- [15] I. Stetcu, C. A. Bertulani, A. Bulgac, P. Magierski, and K. J. Roche, *Phys. Rev. Lett.* **114**, 012701 (2015).
- [16] G. Scamps, C. Simenel, and D. Lacroix, *Phys. Rev. C* **92**, 011602 (2015).
- [17] G. Wlazłowski, K. Sekizawa, P. Magierski, A. Bulgac, and M. M. Forbes, *Phys. Rev. Lett.* **117**, 232701 (2016).
- [18] P. Magierski, K. Sekizawa, and G. Wlazłowski, *Phys. Rev. Lett.* **119**, 042501 (2017).
- [19] G. Scamps and Y. Hashimoto, *Phys. Rev. C* **96**, 031602 (2017).
- [20] V. O. Nesterenko, W. Kleinig, J. Kvasil, P. Vesely, P.-G. Reinhard, and D. S. Dolci, *Phys. Rev. C* **74**, 064306 (2006).
- [21] K. Yoshida and N. V. Giai, *Phys. Rev. C* **78**, 064316 (2008).
- [22] S. Péru and H. Goutte, *Phys. Rev. C* **77**, 044313 (2008).
- [23] D. Pena Arteaga, E. Khan, and P. Ring, *Phys. Rev. C* **79**, 034311 (2009).
- [24] C. Losa, A. Pastore, T. Døssing, E. Vigezzi, and R. A. Broglia, *Phys. Rev. C* **81**, 064307 (2010).

- [25] J. Terasaki and J. Engel, [Phys. Rev. C **82**, 034326 \(2010\)](#).
- [26] K. Yoshida and T. Nakatsukasa, [Phys. Rev. C **83**, 021304 \(2011\)](#).
- [27] K. Yoshida and T. Nakatsukasa, [Phys. Rev. C **88**, 034309 \(2013\)](#).
- [28] J. W. Negele, [Rev. Mod. Phys. **54**, 913 \(1982\)](#).
- [29] S. Levit, [Phys. Rev. C **21**, 1594 \(1980\)](#).
- [30] S. Levit, J. W. Negele, and Z. Paltiel, [Phys. Rev. C **21**, 1603 \(1980\)](#).
- [31] H. Kuratsuji and T. Suzuki, [Phys. Lett. B **92**, 19 \(1980\)](#).
- [32] H. Kuratsuji, [Prog. Theor. Phys. **65**, 224 \(1981\)](#).
- [33] H. Reinhardt, [Nucl. Phys. A **346**, 1 \(1980\)](#).
- [34] D. Bès, R. Broglia, R. Perazzo, and K. Kumar, [Nucl. Phys. A **143**, 1 \(1970\)](#).
- [35] A. Gózdź, K. Pomorski, M. Brack, and E. Werner, [Nucl. Phys. A **442**, 50 \(1985\)](#).
- [36] K. Pomorski, [Int. J. Mod. Phys. E **16**, 237 \(2007\)](#).
- [37] J. Högaasen-Feldman, [Nucl. Phys. **28**, 258 \(1961\)](#).
- [38] M. Cambiaggio, G. Dussel, and M. Saraceno, [Nucl. Phys. A **415**, 70 \(1984\)](#).
- [39] T. Suzuki and Y. Mizobuchi, [Prog. Theor. Phys. **79**, 480 \(1988\)](#).
- [40] R. W. Richardson, [Phys. Lett. **3**, 277 \(1963\)](#).
- [41] R. W. Richardson and N. Sherman, [Nucl. Phys. **52**, 221 \(1964\)](#).
- [42] R. W. Richardson and N. Sherman, [Nucl. Phys. **52**, 353 \(1964\)](#).
- [43] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics, A Course of Theoretical Physics* (Pergamon Press, Oxford, 1965).